

FOURTH ROACH POND TAR12, Piscataquis Co. U.S.G.S. Farrar Mountain, Wadleigh Mountain, Me (7½')

Fishes

Landlocked salmon Brook trout Splake Lake whitefish Rainbow smelt Yellow perch Minnows Lake chub Golden shiner Common shiner Redbelly dace Blacknose dace Creek chub Fallfish White sucker Hornpout (bullhead) Burbot (cusk) Pumpkinseed sunfish Slimy sculpin

Physical Characteristics

Area - 266 acres

Maximum depth - 40 feet

Principal fisheries: Brook trout, landlocked salmon, splake

Fourth Roach Pond can be reached over the network of graveled logging roads leading east from Kokadjo. A well-beaten access trail has developed in recent years.

Suitable brook trout spawning habitat can be found in the lower ends of the tributaries. However, the potential of the pond to sustain a large population of wild brook trout is severely compromised by competition and the lack of dissolved oxygen in the coolest water. Competition increased with the establishment of yellow perch in the 1990s. The brook trout fishery was maintained through the mid-1990s with regular stocking of hatchery-reared brook trout.

Landlocked salmon were stocked in the 1970s in Third Roach Pond. Some moved upstream into Fourth Roach Pond and contributed to its fishery. Salmon stocking in Third Roach Pond resumed in 1997 and those fish are again contributing to the catch in Fourth Roach Pond.

Splake stocking was initiated in 1994. They survive and grow well, taking advantage of the abundance of food. The success of the splake program is probably responsible for the increased interest in the Fourth Roach Pond fishery.

Surveyed - August, 1963 Revised - 2003 Maine Department of Inland Fisheries and Wildlife Funded in part by Federal Aid in Fish Restoration Act under Federal Project F-28-P

L0446E

Temperatures: Surface - 65[°]F 40 feet - 42[°]F