



EXPLANATION

- Syn** Turner Mountain Syenite; coarse-grained, medium to dark red-brown, hornblende-biotite, porphyritic syenite.
- Myl** Mylonite, fine-grained, white to light green, feldspathic.
- Di** Granitic intrusive rocks.
- Dgd** Diorite, medium-grained, dark gray, orthoclase-biotite-pyroxene-quartz.
- EOI** Winterport Granite; foliated granitic rock.
- Bi** Brewer Lake or Stricklen Ridge Intrusive; muscovite-biotite granite with inclusions of Passagassawaukeag Gneiss and Copeland Schist.
- CTC** Roundstone conglomerate, light red.
- CTS** Siltstone and arkosic sandstone, dark red.
- DSV** Vassalboro and Fall Brook Formations; massive, beds 0.3-1.8 m. thick, fine to medium grained, feldspathic wacke, 8-15 cm. thick interbeds of dark gray phyllite, minor black carbonaceous phyllite and feldspathic coarse sand to granule conglomerate.
- Su** Undifferentiated metasediments of probably Silurian age; massive quartzite, metasiltstone, and phyllite.
- Ss** Sangerville Formation; beds 0.1-0.6 m. thick, graded calcareous quartzite with phyllitic tops, phyllitic portions of the beds 5-10 cm. thick, interbedded with dark gray to black phyllitic slate, sometimes rusty weathering.
- Ssp** Maroon and green member; 3-12 mm. thick layers of maroon and green metasiltstone, phyllite, and slate.
- Sw** Waterville Formation; 0.3-1.2 cm. thick layers of phyllite interbedded with 1.2-2.4 cm. thick layers of coarse metasiltstone to very fine quartzite, metasiltstone and quartzite layers exhibit fine internal laminae, grading, and cross-lamination, quartzite layers slightly calcareous.
- SWP** Maroon and green member; 3-12 mm. thick layers of maroon and green metasiltstone, phyllite, and slate.
- Sk** "Kenduskeag Unit"; extremely variable in bedding thickness, sequence of massive quartzite alternating with sequences of 0.6-2.4 cm. thick interbedded phyllite and metasiltstone, portions of the unit consist of sedimentary breccia and chaotic zones of slump origin.
- SKL** Limestone member; 1.2-7 cm. thick layers, silty limestone, interbedded with calcareous siltstone and sandstone.
- Skp** Maroon and green member; 0.3-1.2 cm. thick layers of maroon and green metasiltstone, phyllite, and slate.
- SO** Regularly bedded, 2-4 cm. thick, interbedded phyllitic siltstone and very fine-grained quartzite, quartzite layers have internal laminae that are graded and cross-laminated, gray colored and sericitic.
- Sb(?)** Bucksport Formation; 2-4 cm. thick interlaminated pelite and calcareous silt.
- OSv** Keratophyre, meta-tuff, volcanic breccia, and manganeseiferous siltstone, minor quartzite and black phyllite.
- OSq** Quartz wacke, phyllite, siltstone, and slate.
- OR** Spotted biotite schist, rusty weathering, possibly equivalent to the Cookson Formation to the east.
- Op** Penobscot Formation; rusty weathering, very fine quartzite and black pelite, with black siltstone.
- Oec** Copeland Formation; schist, gray weathering.
- Oer** Rider Bluff Formation; green to gray weathering, thinly laminated, 1-2 cm., thick, non-calcareous, silt-size grains.
- Oev** Green mica schist, meta-volcanic, meta-tuff.
- PEP** Passagassawaukeag Gneiss; interlayered augen gneiss and migmatite.

- Approximate geologic contact
- Fault
- x Outcrop, no bedding or cleavage
- ↘₆₀ Strike and dip of bedding, inclined and vertical. Dot indicates direction of top if determined.
- ↘₈₀ Cleavage, inclined and vertical
- ↘₆₅ Foliation, inclined and vertical



RECONNAISSANCE
BEDROCK GEOLOGY
 OF THE
SAPONAC
QUADRANGLE, MAINE

 BY
JOHN R. GRIFFIN
 1976
 Maine Geological Survey
 DEPARTMENT OF CONSERVATION
 Augusta, Maine 04333
 Walter A. Anderson, State Geologist
OPEN FILE NO. 76-19

