

Surficial Geology

Fayette Quadrangle, Maine

Surficial geologic mapping by
Woodrow B. Thompson (Kennebec and Franklin Cos.)
Geoffrey W. Smith (Androscoggin Co.)

Digital cartography by:
Susan S. Tolman

Robert G. Marvinney
 State Geologist

Cartographic design and editing by:
Robert D. Tucker

Financial assistance provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. 07HQG0079, and by the U.S. Environmental Protection Agency under the provisions of Section 208 of the Federal Water Pollution Control Act, Amendments of 1972.



Maine Geological Survey

Address: 22 State House Station, Augusta, Maine 04333
 Telephone: 207-287-2801 E-mail: mgs@maine.gov
 Home page: <http://www.maine.gov/doc/trinc/trinc.htm>

Open-File No. 08-44

2008
 This map supersedes
 Open-File Map 77-42.

SURFICIAL GEOLOGY OF MAINE

Continental glaciers like the ice sheet now covering Antarctica probably extended across Maine several times during the Pleistocene Epoch, between about 1.5 million and 10,000 years ago. The slow-moving ice superficially changed the landscape as it scraped over mountains and valleys (Figure 1), eroding and transporting boulders and other rock debris for miles (Figure 2). The sediments that cover much of Maine are largely the product of glaciation. Glacial ice deposited some of these materials, while others were washed into the sea or accumulated in meltwater streams and lakes as the ice receded. Earlier stream patterns were disrupted, creating hundreds of ponds and lakes across the state. The map at left shows the pattern of glacial sediments in the Fayette quadrangle.

The most recent "Ice Age" in Maine began about 30,000 years ago when an ice sheet spread southward over New England (Stone and Borns, 1986). During its peak, the ice was several thousand feet thick and covered the highest mountains in the state. The weight of this huge glacier actually caused the land surface to sink hundreds of feet. Rock debris frozen into the base of the glacier abraded the bedrock surface over which the ice flowed. The grooves and fine scratches (striations) resulting from this scraping process are often seen on freshly exposed bedrock, and they are important indicators of the direction of ice movement (Figure 3). Erosion and sediment deposition by the ice sheet combined to give a streamlined shape to many hills, with their long dimension parallel to the direction of ice flow. Some of these hills (drumlins) are composed of dense glacial sediment (till) plastered under great pressure beneath the ice.

A warming climate forced the ice sheet to start receding as early as 21,000 calendar years ago, soon after it reached its southernmost position on Long Island (Ridge, 2004). The edge of the glacier withdrew from the continental shelf east of Long Island and reached the present position of the Maine coast by about 16,000 years ago (Borns and others, 2004). Even though the weight of the ice was removed from the land surface, the Earth's crust did not immediately spring back to its normal level. As a result, the sea flooded much of southern Maine as the glacier retreated to the northwest. Ocean waters extended far up the Kennebec and Penobscot valleys, reaching present elevations of up to 420 feet in the central part of the state.

Great quantities of sediment washed out of the melting ice and into the sea, which was in contact with the receding glacier margin. Sand and gravel accumulated as deltas (Figure 4) and submarine fans where streams discharged along the ice front, while the finer silt and clay dispersed across the ocean floor. The shells of clams, mussels, and other invertebrates are found in the glacial-marine clay that blankets lowland areas of southern Maine. Ages of these fossils tell us that ocean waters covered parts of Maine until about 13,000 years ago. The land rebounded as the weight of the ice sheet was removed, forcing the sea to retreat.


References Cited

Borns, H. W., Jr., Doner, L. A., Dorion, C. C., Jacobson, G. L., Jr., Kaplan, M. R., Kreutz, K. J., Lowell, T. V., Thompson, W. B., and Weddle, T. K., 2004. The deglaciation of Maine, U.S.A., in Ehlers, J., and Gibbard, P. L., eds., *Quaternary Glaciations - Extent and Chronology, Part II: North America*, Amsterdam, Elsevier, p. 89-109.

Davis, R. B., and Jacobson, G. L., Jr., 1985. Late-glacial and early Holocene landscapes in northern New England and adjacent areas of Canada. *Quaternary Research*, v. 23, p. 341-368.


Ridge, J. C., 2004. The Quaternary glacial history of western New England with correlations to surrounding areas, in Ehlers, J., and Gibbard, P. L., eds., *Quaternary Glaciations - Extent and Chronology, Part II: North America*, Amsterdam, Elsevier, p. 169-199.

Stone, B. D., and Borns, H. W., Jr., 1986. Pleistocene glacial and interglacial stratigraphy of New England, Long Island, and adjacent Georges Bank and Gulf of Maine, in Shrivya, V., Bowen, D. O., and Richmond, G. M. (editors), *Quaternary glaciations in the northern hemisphere: Quaternary Science Reviews*, v. 5, p. 39-52.



SOURCES OF INFORMATION

Editing and updating of map units completed by Woodrow B. Thompson in 2008 for the STATEMAP program. Original reconnaissance field mapping by G. W. Smith and W. B. Thompson in the mid 1970's, supplemented by later field checks.



Topographic base from U.S. Geological Survey Fayette quadrangle, scale 1:24,000 using standard U.S. Geological Survey topographic map symbols.

The use of industry, firm, or local government names on this map is for location purposes only and does not impure responsibility for any present or potential effects on the natural resources.

- Hw** Wetland deposits - Peat, muck, silt, and clay. Deposited by accumulation of organic-rich sediments in poorly drained areas on valley floors. Until may grade into or include areas of stream alluvium.
- Oe** Eolian deposits - Windblown sand derived from glacial-lake sediments in the Androscoggin River basin. Includes longitudinal dunes oriented parallel to the prevailing wind direction when the dunes formed. Smaller unmapped areas of eolian sand may occur elsewhere in the quadrangle.
- Pm** Glaciomarine sediments, undifferentiated - Sand, gravel, and clay-silt deposited in the late-glacial sea. May include deltaic and submarine fan deposits formed near the glacier margin and locally modified by postglacial erosion.
- Pp** Presumpscot Formation - Glaciomarine silt, clay, and sand deposited on the late-glacial sea floor.
- Pg** Glacial meltwater deposits - Sand and gravel deposited by glacial meltwater.
- Pge** Esker deposits - Ridges of sand and gravel deposited by glacial meltwater streams in subglacial tunnels.
- Pt** Till - Loose to very compact, poorly sorted, massive to weakly stratified mixture of sand, silt, and gravel-size rock debris deposited by glacial ice. Locally includes lenses of waterlaid sand and gravel.
- Bedrock outcrops / thin-drift areas** - Ruled pattern indicates areas where outcrops are common and/or surficial sediments are generally less than 10 ft thick (mapped partly from air photos). Dots show individual outcrops.
- Contact** - Boundary between map units. Dashed where approximately located.
- Grooved till surface** - Narrow ridges and grooves in till sculpted by flow of glacial ice.
- Glacially streamlined hill** - Symbol shows trend of long axis of hill, which is parallel to former glacial ice-flow direction.
- Glacial striation locality** - Arrow shows ice-flow direction inferred from striations on bedrock. Dot marks point of observation. Number is azimuth (in degrees) of flow direction.
- K** Kettle - Depression created by melting of buried glacial ice and collapse of overlying sediments.
- Meltwater channel** - Channel eroded by glacial meltwater stream. Arrow shows inferred direction of water flow.
- Crest of esker** - Chevrons show trend of esker ridge and point in direction of glacial meltwater flow.
- Area of large boulders** - Area where many boulders are scattered across the ground surface. Shown only where observed, so these areas probably are more extensive than indicated on the map.

USES OF SURFICIAL GEOLOGY MAPS

A surficial geology map shows all the loose materials such as till (commonly called hardpan), sand and gravel, or clay, which overlie solid ledge (bedrock). Bedrock outcrops and areas of abundant bedrock outcrops are shown on the map, but varieties of the bedrock are not distinguished (refer to bedrock geology map). Most of the surficial materials are deposits formed by glacial and deglacial processes during the last stage of continental glaciation, which began about 25,000 years ago. The remainder of the surficial deposits are the products of postglacial geologic processes, such as river floodplains, or are attributed to human activity, such as fill or other land-modifying features.

The map shows the areal distribution of the different types of glacial features, deposits, and landforms as described in the map explanation. Features such as striations and moraines can be used to reconstruct the movement and position of the glacier and its margin, especially as the ice sheet melted. Other ancient features include shorelines and deposits of glacial lakes or the glacial sea, now long gone from the state.

This glacial geology history of the quadrangle is useful to the larger understanding of past earth climate, and how our region of the world underwent recent geologically significant climatic and environmental changes. We may then be able to use this knowledge in anticipation of future similar changes for long-term planning efforts, such as coastal development or waste disposal.

Surficial geology maps are often best used in conjunction with related maps such as surficial materials maps or significant sand and gravel aquifer maps for anyone wanting to know what lies beneath the land surface. For example, these maps may aid in the search for water supplies, or economically important deposits such as sand and gravel for aggregate or clay for bricks or pottery. Environmental issues such as the location of a suitable landfill site or the possible spread of contaminants are directly related to surficial geology. Construction projects such as locating new roads, excavating foundations, or siting new homes may be better planned with a good knowledge of the surficial geology of the site. Refer to the list of related publications below.

OTHER SOURCES OF INFORMATION

- Locke, D. B., 2008. Surficial materials of the Fayette quadrangle, Maine: Maine Geological Survey, Open-File Map 08-65.
- Neil, C. D. (compiler), 1998. Significant sand and gravel aquifers in the Fayette quadrangle, Maine: Maine Geological Survey, Open-File Map 98-219.
- Thompson, W. B., 1979. Surficial geology handbook for coastal Maine: Maine Geological Survey, 68 p. (out of print).
- Thompson, W. B., and Borns, H. W., Jr., 1985. Surficial geologic map of Maine: Maine Geological Survey, scale 1:500,000.



Figure 1: "The Bubbles" and Jordan Pond in Acadia National Park. These hills and valleys were sculpted by glacial erosion. The pond was dammed behind a moraine ridge during retreat of the ice sheet.



Figure 2: Dagget's Rock in Phillips. This is the largest known glacially transported boulder in Maine. It is about 100 feet long and estimated to weigh 8,000 tons.




Figure 3: Granite ledge in Westbrook, showing polished and grooved surface resulting from glacial abrasion. The grooves and shape of the ledge indicate ice flow toward the southeast.




Figure 4: Glaciomarine delta in Franklin, formed by sand and gravel washing into the ocean from the glacier margin. The flat delta top marks approximate former sea level. Kettle hole in foreground was left by retreating glacier margin.




Figure 5: Esker cutting across Kezar Five Ponds, Watford. The ridge consists of sand and gravel deposited by meltwater flowing in a tunnel beneath glacial ice.




Figure 6: Aerial view of moraine ridges in blueberry field, Sedgwick (note dirt road in upper right for scale). Each bouldery ridge marks a position of the retreating glacier margin. The ice receded from right to left.




Figure 7: Sand dune in Wayne. This and other "deserts" in Maine formed as windblown in late-glacial time blew sand out of valleys, often depositing it as dune fields on hillsides downwind. Some dunes were reactivated in historical times when grazing animals stripped the vegetation cover.



Figure 8: Songo River delta and Songo Beach, Sebago Lake State Park, Naples. These deposits are a typical glacial feature formed in Maine since the Ice Age.