

# Surficial Geology

# Gardiner Quadrangle, Maine

Surficial geologic mapping by  
**Woodrow B. Thompson**

Digital cartography by  
**Robert A. Johnston**

**Robert G. Marvinney**  
State Geologist

Cartographic design and editing by  
**Robert D. Tucker**

Funding for the preparation of this map was provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. 04HQG0005.



**Maine Geological Survey**  
Address: 22 State House Station, Augusta, Maine 04333  
Telephone: 207-287-2801 E-mail: mgs@maine.gov  
Home page: http://www.maine.gov/doc/nrmc/nrmc.htm

**Open-File No. 09-8**  
**2009**  
This map supersedes  
Open-File Map 07-103.

## SURFICIAL GEOLOGY OF MAINE

Continental glaciers like the ice sheet now covering Antarctica probably extended across Maine several times during the Pleistocene Epoch, between about 1.5 million and 10,000 years ago. The slow-moving ice superficially changed the landscape as it scraped over mountains and valleys, eroding and transporting boulders and other rock debris for miles. The sediments that cover much of Maine are largely the product of glaciation. Glacial ice deposited some of these materials, while others washed into the sea or accumulated in meltwater streams and lakes as the ice receded. Earlier stream patterns were disrupted, creating hundreds of ponds and lakes across the state. The map at left shows the pattern of glacial sediments in the Gardiner quadrangle.

The most recent "Ice Age" in Maine began about 25,000 years ago when an ice sheet spread southward over New England (Stone and Borns, 1986). During its peak, the ice was several thousand feet thick and covered the highest mountains in the state. The weight of this huge glacier actually caused the land surface to sink hundreds of feet. Rock debris frozen into the base of the glacier abraded the bedrock surface over which the ice flowed. The grooves and fine scratches (striations) resulting from this scraping process are often seen on freshly exposed bedrock, and they are important indicators of the direction of ice movement. Erosion and sediment deposition by the ice combined to give a streamlined shape to many hills, with their long dimension parallel to the direction of ice flow. Some of these hills (drumlins) are composed of dense glacial sediment (till) plastered under great pressure beneath the ice.

A warming climate forced the ice sheet to start retreating as early as 21,000 years ago, soon after it reached its southernmost position on Long Island (Sarkin, 1986). The edge of the glacier withdrew from the continental shelf east of Long Island and reached the present position of the Maine coast by 13,800 years ago (Dorion, 1993). Even though the weight of the ice was removed from the land surface, the Earth's crust did not immediately spring back to its normal level. As a result, the sea flooded much of southern Maine as the glacier retreated to the northwest. Ocean waters extended far up the Kennebec and Penobscot valleys, reaching present elevations of up to 420 feet in the central part of the state.

Great quantities of sediment washed out of the melting ice and into the sea, which was in contact with the retreating glacier margin. Sand and gravel accumulated as deltas and submarine fans where streams discharged along the ice front, while the finer silt and clay dispersed across the ocean floor. The shells of clams, mussels, and other invertebrates are found in the glacial-marine clay that blankets low land areas of southern Maine. Age dates on these fossils tell us that ocean waters covered parts of Maine until about 11,000 years ago, when the land surface rebounded as the weight of the ice sheet was removed.


Meltwater streams deposited sand and gravel in tunnels within the ice. These deposits remained as ridges (eskers) when the surrounding ice disappeared. Maine's esker systems can be traced for up to 100 miles, and are among the longest in the country.

Other sand and gravel deposits formed as mounds (kames) and terraces adjacent to melting ice, or as outwash in valleys in front of the glacier. Many of these water-laid deposits are well layered, in contrast to the chaotic mixture of boulders and sediment of all sizes (till) that was released from dirty ice without subsequent reworking. Ridges consisting of till or washed sediments (moraines) were constructed along the ice margin in places where the glacier was still actively flowing and covering rock debris to its terminus. Moraine ridges are abundant in the zone of former marine submergence, where they are useful indicators of the pattern of ice retreat.

The last remnants of glacial ice probably were gone from Maine by 10,000 years ago. Large sand dunes accumulated in late-glacial time as winds picked up outwash sand and blew it onto the east sides of river valleys, such as the Androscoggin and Saco valleys. The modern stream network became established soon after deglaciation, and organic deposits began to form in peat bogs, marshes, and swamps. Tundra vegetation bordering the ice sheet was replaced by changing forest communities as the climate warmed (Davis and Jacobson, 1985). Geologic processes are by no means dormant today, however, since rivers and wave action modify the land, and worldwide sea level is gradually rising against Maine's coast.


## References Cited

- Davis, R. B., and Jacobson, G. L., Jr., 1985. Late-glacial and early Holocene landscapes in northern New England and adjacent areas of Canada: *Quaternary Research*, v. 23, p. 341-368.
- Dorion, C. C., 1993. A chronology of deglaciation and accompanying marine transgression in Maine. *Geological Society of America, Abstracts with Programs*, v. 25, no. 2, p. 12.
- Sarkin, L., 1986. Pleistocene stratigraphy of Long Island, New York. In Caldwell, D. W. (editor), *The Wisconsin stage of the first geological district, eastern New York*. New York State Museum, Bull. 455, p. 6-21.
- Stone, B. D., and Borns, H. W., Jr., 1986. Pleistocene glacial and interglacial stratigraphy of New England, Long Island, and adjacent Georges Bank and Gulf of Maine. In Sibrava, V., Bowen, D. Q., and Richmond, G. M. (editors), *Quaternary glaciations in the northern hemisphere: Quaternary Science Reviews*, v. 5, p. 39-52.



### SOURCES OF INFORMATION

Surficial geologic mapping of the Gardiner quadrangle was conducted by Woodrow B. Thompson in 2004-05 for the STATEMAP program and modified by 2008 field data. Some of the data included here were collected by W. B. Thompson during reconnaissance surficial mapping of the Gardiner 15-minute quadrangle in 1975 and scattered observations during the 1980's and 1990's.



Topographic base from U.S. Geological Survey Gardiner quadrangle, scale 1:24,000 using standard U.S. Geological Survey topographic map symbols.

The use of industry, firm, or local government names on this map is for location purposes only and does not implicate responsibility for any present or potential effects on the natural resources.

- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                              |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Ha</b> Stream alluvium - Sand, gravel, and silt deposited on flood plains of the Kennebec River and other streams. May include some wetland deposits.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <b>Pge</b> Esker - Sand and gravel deposited by glacial meltwater stream in a tunnel beneath the ice. Chevron symbols show inferred direction of former stream flow.                                                                                                                                                                         |
| <b>Hw</b> Wetland deposits - Peat, muck, silt, and clay in poorly drained areas.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | <b>Pt</b> Till - Loose to very compact, poorly sorted, massive to weakly stratified mixture of sand, silt, and gravel-size rock debris deposited by glacial ice. Locally includes lenses of water-laid sand and gravel. Boulders commonly present on ground surface.                                                                         |
| <b>Qst</b> Stream terraces - Sand and gravel deposited by the Kennebec River at elevations higher than the most recent flood plains.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>Bedrock outcrops/thin-drift areas</b> - Ruled pattern indicates areas where bedrock outcrops are common and/or surficial sediments are generally less than 10 ft thick. Mapped from air photos and ground observations. Actual thin-drift areas probably are more extensive than shown. Dots mark locations of small individual outcrops. |
| <b>Pmn</b> Marine nearshore deposits - Sandy to gravelly sediments formed when marine processes reworked older glacial deposits during recession of the sea.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | <b>af</b> Artificial fill - Variable mixtures of earth, rock, and/or man-made materials used as fill for roads. Shown only where large enough to affect the contour pattern on the topographic map.                                                                                                                                          |
| <b>Pp</b> Presumpscot Formation - Glaciomarine silt, clay, and sand deposited on the late-glacial sea floor.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | <b>Contact</b> - Boundary between map units, dashed where location is approximate.                                                                                                                                                                                                                                                           |
| <b>Pmf</b> Glaciomarine fan - Sand and gravel deposited as a submarine fan at the glacier margin during recession of the late Wisconsinan ice sheet in the Kennebec River valley.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | <b>Moraine ridge</b> - Line shows inferred crest of moraine ridge deposited along the retreating margin of the most recent (late Wisconsinan) glacial ice sheet.                                                                                                                                                                             |
| <b>Pp/Pgi</b> Presumpscot Formation overlying ice-contact deposits - Sand and gravel (Pgi) deposited as eskers and/or glaciomarine fans adjacent to glacial ice in the Kennebec River valley. The ice-contact deposits are overlain by variable thicknesses of glaciomarine silt, clay, and sand (Pp). These units were not mapped individually because of poor exposure and complex stratigraphy.                                                                                                                                                                                                                                                                                                                                                                                                                      | <b>Glacially streamlined hill</b> - Symbol shows long axis of hill or ridge shaped by flow of glacial ice, and which is parallel to former ice-flow direction.                                                                                                                                                                               |
| <b>Pp/Pm/Pg</b> Presumpscot Formation overlying glaciomarine fan and esker deposits. These deposits form a complex assemblage in the Kennebec River valley. A ridge of coarse esker gravel (Pge) in the valley bottom is generally buried by submarine fan deposits (Pmf) comprised of stratified sand and gravel. Variable thicknesses of glaciomarine silt, clay, and sand (Pp) overlie the sand and gravel units, and are locally overlain in turn by coarser sediments formed by the wave and current action as relative sea level fell. These units could not be distinguished accurately at the scale of the map, due to their complex interrelations and limited fresh exposures. Bedrock has been exposed in some of the deeper gravel pits, and large portions of the sand and gravel units have been removed. | <b>Glacial striation locality</b> - Arrow shows ice-flow direction inferred from striations on bedrock. Dot marks point of observation. Number is azimuth (in degrees) of flow direction. Flagged trend is older.                                                                                                                            |
| <b>Pgi</b> Ice-contact deposits. Miscellaneous sand and gravel deposits formed in contact with glacial ice.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | <b>Crest of esker</b> - Alignment of symbols shows trend of esker ridge. Chevrons point in direction of meltwater flow.                                                                                                                                                                                                                      |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | <b>Marine fossil locality</b> - Age in radiocarbon years.                                                                                                                                                                                                                                                                                    |

### USES OF SURFICIAL GEOLOGY MAPS

A surficial geology map shows all the loose materials such as till (commonly called hardpan), sand and gravel, or clay, which overlie solid ledge (bedrock). Bedrock outcrops and areas of abundant bedrock outcrops are shown on the map, but varieties of the bedrock are not distinguished (refer to bedrock geology map). Most of the surficial materials are deposits formed by glacial and deglacial processes during the last stage of continental glaciation, which began about 25,000 years ago. The remainder of the surficial deposits are the products of postglacial geologic processes, such as river floodplains, or are attributed to human activity, such as fill or other land-modifying features.

The map shows the areal distribution of the different types of glacial features, deposits, and landforms as described in the map explanation. Features such as striations and moraines can be used to reconstruct the movement and position of the glacier and its margin, especially as the ice sheet melted. Other ancient features include shorelines and deposits of glacial lakes of the glacial sea, now long gone from the state. This glacial geologic history of the quadrangle is useful to the larger understanding of past earth climate, and how our region of the world underwent recent geologically significant climatic and environmental changes. We may then be able to use this knowledge in anticipation of future similar changes for long-term planning efforts, such as coastal development or waste disposal.

Surficial geology maps are often best used in conjunction with related maps such as surficial materials maps or significant sand and gravel aquifer maps for any water wanting to know what lies beneath the land surface. For example, these maps may aid in the search for water supplies, or economically important deposits such as sand and gravel for aggregate or clay for bricks or pottery. Environmental issues such as the location of a suitable landfill site or the possible spread of contaminants are directly related to surficial geology. Construction projects such as locating new roads, excavating foundations, or siting new homes may be better planned with a good knowledge of the surficial geology of the site. Refer to the list of related publications below.

### OTHER SOURCES OF INFORMATION

- Thompson, W. B., and Locke, D. B., 2005. Surficial materials of the Gardiner quadrangle, Maine. Maine Geological Survey, Open-File Map 05-2.
- Neil, C. D., 1999. Significant sand and gravel aquifers of the Gardiner quadrangle, Maine. Maine Geological Survey, Open-File Map 99-36.
- Thompson, W. B., 1979. Surficial geology handbook for coastal Maine. Maine Geological Survey, 68 p. (out of print).
- Thompson, W. B., and Borns, H. W., Jr., 1985. Surficial geologic map of Maine. Maine Geological Survey, scale 1:500,000.



Figure 1: Glacially abraded granite outcrop near intersection of Miles Lane and Maloy Avenue, north of Route 9/126 in West Gardiner. The asymmetric profile of the outcrop, with the gentler slope facing "up-glacier," and grooves on the ledge surface show that the glacier flowed east-southeast (right to left, as seen in photo).




Figure 2: Glacial grooves on top surface of gneiss outcrop, west side of Kennebec River in South Gardiner. Pencil is parallel to ice flow direction (129°).



Figure 3: Pit exposure north of Beedle Road in Richmond, showing glacial till (above shovel) overlying and interlayered with deformed sand beds. The till was deposited at the base of the ice when it readvanced a short distance over the northern margin of a submarine fan.



Figure 4: View looking south-southeast across submarine fan in Richmond (same pit as Figure 3). The fan is composed of sand and gravel that washed into the sea at the edge of the last glacial ice sheet during its retreat from the area. A thin deposit of gray glacial-marine clay overlies the fan near left edge of photo.



Figure 5: Coarse esker gravel on west side of Kennebec River in Gardiner and South Gardiner. The esker was deposited by a meltwater stream in a subglacial ice tunnel. It is part of a long esker system that follows the Kennebec Valley, but much of this deposit is concealed under glacial-marine clay in the Gardiner area.



Figure 6: Gravel pit on east side of Kennebec River in Pittston, showing cross-section of esker ridge. The esker is composed of gravel, and it is directly overlain by gray glacial-marine clay. Unlike most pits in the Kennebec Valley, this exposure does not show a submarine fan deposit between the esker and younger clay unit.




Figure 7: Glacial-marine clay (Presumpscot Formation) formerly exposed at construction site in Gardiner, near west end of Kennebec River bridge. Lower part of section shows fresh "blue clay," while the upper part has been oxidized to a brownish color.




Figure 8: Pit face on east side of Kennebec River in Pittston, showing glacial-marine clay (Presumpscot Formation) overlying sand and gravel (submarine fan). The latter unit was eroded by meltwater currents or a submarine landslide that truncated the fan beds prior to deposition of the clay.