



Department of Health and Human Services  
Maine Center for Disease Control and Prevention  
286 Water Street  
11 State House Station  
Augusta, Maine 04333-0011  
Tel: (207) 287-8016; Fax (207) 287-9058  
TTY Users: Dial 711 (Maine Relay)

## Maine Health Alert Network (HAN) System

### PUBLIC HEALTH ADVISORY

**To:** Health Care Providers  
**From:** Dr. Isaac Benowitz, State Epidemiologist  
**Subject:** U.S. CDC: First Case of Clade I Mpox Diagnosed in the United States  
**Date / Time:** Tuesday, November 26, 2024, at 4:30PM  
**Priority:** Normal  
**Pages:** 6  
**Message ID:** 2024PHADV040

Please take a moment to review this information from U.S. CDC on the first detection of clade I mpox in the United States. Maine has not detected any cases of clade I mpox.

Information for health care providers on mpox testing, vaccination, treatment, and infection control can be found at <https://www.maine.gov/dhhs/mecdc/infectious-disease/epi/zoonotic/monkeypox-providers.shtml>.

Health care providers who suspect mpox in a person with travel to Democratic Republic of the Congo (DRC) or neighboring countries should send specimens to Maine's Health and Environmental Testing Laboratory (HETL) for mpox testing. Results for mpox testing conducted at HETL are generally available within two business days.

For patients without travel to DRC and neighboring countries, Maine CDC encourages clinicians to use commercial laboratories for mpox testing.

Travelers to DRC or other countries with sustained spread of clade I mpox, regardless of sexual orientation or gender identity, should be made aware of activities associated with cases and should be vaccinated with two doses of JYNNEOS if they anticipate certain sexual exposures while traveling. To find an mpox vaccination provider near you, visit <https://mpoxvaccine.cdc.gov/>.

Maine CDC strongly encourages clinicians evaluating persons for mpox to also evaluate and screen for sexually transmitted infections (STIs), including HIV, syphilis, gonorrhea, and chlamydia, per [U.S. CDC STI Treatment Guidelines](#). Maine CDC currently recommends HIV testing at least every 3 months for persons with ongoing risk factors. This [interim recommendation](#) is related to the HIV cluster in Penobscot County.

Any confirmed case of mpox should be reported to Maine CDC by electronic laboratory report or phone. To contact Maine CDC, please call the 24/7 disease reporting number at 800-821-5821.

## U.S. CDC: First Case of Clade I Mpox Diagnosed in the United States

### Summary

The Centers for Disease Control and Prevention (U.S. CDC) is issuing this Health Alert Network (HAN) Health Advisory to provide information about the first case of clade I mpox diagnosed in the United States and recommendations to clinicians about preventing, diagnosing, treating, and reporting mpox cases. On November 15, 2024, the California Department of Public Health (CDPH) confirmed the [first reported case of clade I mpox](#) in

the United States. This individual had recently traveled to areas experiencing clade I monkeypox virus (MPXV) transmission and sought medical care for mpox symptoms in the United States. Consistent with other recent clade I mpox cases, the patient has relatively mild illness and is recovering. U.S. CDC and the local and state health departments are investigating potential contacts; no additional cases in the United States have been detected as of November 18, 2024. The [risk of clade I mpox to the public](#) in the United States remains low.

Since March 2024, U.S. CDC has been working with local, tribal, state, and territorial public health authorities to prepare for potential cases of clade I mpox in the United States by enhancing surveillance, detection, and reporting capacities of existing domestic public health systems and structures. This reported case demonstrates that these systems are working as intended. There is no change to U.S. CDC clinical or travel guidance on clade I mpox since [HAN Health Update 516](#). Clinicians should be aware of [mpox symptoms](#), ask patients with comparable signs and symptoms about recent [travel history and other risk factors](#) for mpox, and [consider MPXV testing](#). Given the widespread outbreaks in Central and Eastern Africa, additional travel-associated cases may be reported in the future in the United States. [Suspected and confirmed cases](#) of clade I mpox should be reported to **Maine CDC** as soon as possible. This includes *orthopoxvirus* generic (i.e., non-variola *orthopoxvirus*) **positive** and clade II **negative** test results from a patient with travel history to country affected by clade I mpox. U.S. CDC recommends vaccination to people who are [eligible for mpox vaccine](#), including those who may have a [recent MPXV exposure](#).

## Background

MPXV has two distinct genetic clades: clade I (with subclades Ia and Ib) is [endemic to some countries in Central Africa](#), and clade II (with subclades IIa and IIb) is historically [endemic to some countries in West Africa](#). MPXV transmission in countries where the virus is endemic typically occurs via exposure to infected wildlife with subsequent [person-to-person spread](#) via close contact (including intimate, sexual, or household contact) with a person with mpox, or direct contact with infectious respiratory secretions (e.g., snot, mucus) or contaminated objects (e.g., bedding). Clade I and [clade II](#) mpox present similarly, and, as with clade II mpox, [clinical management](#) of clade I mpox is based on the severity of illness at diagnosis and the [potential for severe or prolonged mpox](#).

From January 1 through November 15, 2024, about [12,000 confirmed cases of clade I mpox](#) and at least 47 deaths have been reported in [Central and Eastern African countries](#). These countries include Burundi, Central African Republic, Democratic Republic of the Congo, Republic of the Congo, Rwanda, and Uganda. Data from affected countries indicate that a large proportion of clade I mpox cases among adults were associated with [heterosexual contact](#). Transmission to close contacts within households, including to children, also has been reported.

Travel-associated clade I mpox cases have been reported in Germany (1), India (1), Kenya (17), Sweden (1), Thailand (1), the United Kingdom (UK) (4), Zambia (1), and Zimbabwe (2) so far in 2024, and no onward spread has been reported except to close household contacts in Kenya and the UK. Current data suggest that subclade Ib may be less severe. Clade Ib mpox has a lower death rate (less than 1%) than clade Ia both in and outside of Africa. No deaths have occurred in travel-associated clade Ib mpox cases in countries outside of Africa; for a subset of these cases for which clinical data are available, relatively mild disease courses were described.

On November 15, 2024, CDPH confirmed through laboratory testing the first reported case of clade I mpox in the United States. The case was diagnosed in a person who recently visited an area with a clade I mpox outbreak. Based on the patient's travel history and symptoms, clinical specimens were tested; PCR was positive for non-variola *orthopoxvirus* and negative by PCR for clade II. Subsequent PCR testing for clade I mpox was positive. Specimens have been sent to U.S. CDC for additional virus characterization.

The individual received care in the United States and is isolating from others. The patient, who has no underlying health conditions, has not had any severe manifestations of disease, and symptoms are improving. U.S. CDC is working closely with the local and state health authorities to rapidly investigate the circumstances surrounding this case and to prevent spread of the virus. As of November 18, no additional clade I mpox cases have been reported in the United States.

Since March 2024, U.S. CDC has been working with local, tribal, state, and territorial public health partners and other U.S. Government agencies, to prepare for potential cases of clade I mpox in the United States by enhancing surveillance, detection, and reporting capacities of existing public health systems and structures. This reported

case demonstrates that these systems are working as intended. U.S. CDC guidance for [clinical care](#), [prevention](#), [vaccination](#), [infection prevention and control](#), and exposure risks in [community](#), [healthcare](#), and [travel](#) settings have not changed. Guidance for travelers is unchanged from that described in [HAN Health Update 516](#); see also [HAN Health Update 513](#) and [HAN Health Advisory 501](#). The [overall risk of clade I mpox to the public in the United States remains low](#).

## Recommendations for Clinicians and Public Health Practitioners

### Evaluation and Diagnosis

- **Consider mpox as a possible diagnosis in patients with [epidemiologic characteristics](#) and [lesions or other clinical signs and symptoms](#) consistent with mpox.**
  - This includes symptomatic people who have been in [Central or Eastern Africa](#) (including, but not limited to, Burundi, Central African Republic, Democratic Republic of the Congo, Kenya, Republic of the Congo, Rwanda, Uganda, Zambia, or Zimbabwe) in the previous 21 days.
  - This also includes people who had close or intimate contact with symptomatic people who have been in these countries.
  - An up-to-date list of [countries affected by clade I mpox outbreaks](#) is available on the U.S. CDC website.
- Follow U.S. CDC guidance on mpox [infection prevention and control](#) to minimize transmission risk when evaluating and providing care to patients with suspected mpox.
- Ask patients with signs and symptoms of mpox but no recent travel whether they have had contact with people who had recently been in [Central or Eastern Africa](#) and who were symptomatic for mpox.
- Consider mpox as a possible diagnosis if a clinically consistent presentation occurs, even in people vaccinated for or [previously diagnosed with mpox](#).
- Advise all patients suspected of having mpox to stay at home and [isolate themselves](#) from others until mpox has been ruled out by laboratory testing. In the event of a positive mpox diagnosis, advise patients to isolate until their mpox lesions have cleared up and fresh skin has formed, which could take several weeks.
- Test all [suspected cases](#) for MPXV. If a symptomatic patient reports travel to [Central or Eastern Africa](#) in the 21 days prior to relevant symptom onset, work with Maine CDC to facilitate testing for MPXV that includes clade I MPXV testing. In most situations, specimens should be sent to the appropriate state public health laboratory or a commercial laboratory for initial testing. If you are authorized by Maine CDC to send specimens directly to U.S. CDC for testing, contact U.S. CDC at [poxviruslab@cdc.gov](mailto:poxviruslab@cdc.gov) for information about specimen types accepted, labeling, specimen storage, and shipping timeframes.
- Follow [specimen collection guidelines](#) (including collecting two swabs per 2-3 lesions) to ensure specimen availability for clade-specific testing. This testing will help distinguish cases that are part of the [ongoing clade II mpox global outbreak](#) from those that are part of this clade I outbreak.
  - Avoid unroofing or aspirating lesions (or otherwise using sharp instruments for mpox testing) to minimize the risk of a sharps injury.
- Send clinical specimens to a laboratory that can perform clade-specific MPXV testing **as quickly as possible**. If you need assistance locating relevant laboratories in your area, email [poxvirus@cdc.gov](mailto:poxvirus@cdc.gov).
- Promptly report suspected cases of clade I mpox to Maine CDC.
- U.S. CDC encourages the state health department and diagnosing clinician to contact the U.S. CDC Emergency Operations Center (EOC) at 770-488-7100 and request a clinical mpox consult after clade I mpox is diagnosed, regardless of the severity of illness.

### Treatment

- Promptly consult Maine CDC or U.S. CDC ([poxvirus@cdc.gov](mailto:poxvirus@cdc.gov)) about any mpox cases for which severe manifestations might occur (e.g., in people with advanced HIV infection or severe immunocompromise).
- Inform all patients, including those with mild disease, about the [STOMP Trial](#) and encourage to consider enrollment. To enroll in STOMP, call 1-855-876-9997.
- For patients who are not eligible for inclusion in the STOMP trial and who meet U.S. CDC's [expanded use Investigational New Drug \(EA-IND\) eligibility](#) for tecovirimat treatment, contact Maine CDC to see if oral tecovirimat remains available from prior prepositioned supplies; they will facilitate consultation with U.S. CDC ([poxvirus@cdc.gov](mailto:poxvirus@cdc.gov)).

## Prevention

- Recommend vaccination to people who are [eligible for mpox vaccine](#), including those who may have a [recent MPXV exposure](#).
- Continue to follow U.S. CDC's [current vaccine guidance](#) to prevent mpox.
  - Two doses of JYNNEOS vaccine [offer substantial protection against mpox](#), and are expected to offer protection regardless of clade.
  - If people at risk for mpox have only received one dose, remind them to get a second dose as soon as possible.
  - More than two JYNNEOS vaccine doses ("boosters") are not currently recommended.
- Discuss mpox prevention and risk reduction strategies with all travelers to countries with ongoing human-to-human transmission of clade I MPXV. An [updated list of the countries](#) with ongoing spread of clade I MPXV is available on the U.S. CDC website.
- Discuss patients' [sexual history](#) and travel plans, including if patients anticipate sexual or intimate activity during travel.
- Advise patients that mpox exposure risk is often associated with sexual or intimate contact.
- Remind patients that mpox is not spread through casual contact, such as someone might have in public spaces like markets, offices, classrooms, public transit, or air travel.
- Counsel patients on [activities that may increase risk](#) for MPXV exposure and risk reduction strategies if they have plans to travel to a [country where ongoing human-to-human transmission](#) of clade I MPXV is occurring. Travelers to affected countries should:
  - Avoid close contact with people who are sick with [signs and symptoms](#) of mpox, including skin or genital lesions.
  - Avoid contact with contaminated materials used by people who are sick, such as clothing, bedding, toothbrushes, sex toys, or materials used in healthcare settings.
  - Avoid contact with animals that can carry the virus that causes mpox or their products (e.g., bushmeat, lotions, hides) in areas where mpox is endemic, particularly in Central or West Africa.
- Clinicians should counsel patients about [what to do to prevent household transmission if they have mpox symptoms](#), including staying away from other people, not sharing things they have touched with others, and cleaning and disinfecting the spaces they occupy regularly to limit household contamination.

## Recommendations for Laboratories

- According to [Advisory Committee on Immunization Practices \(ACIP\)](#) recommendations, employers should offer pre-exposure *orthopoxvirus* vaccination to workers at risk of occupational exposure. Two vaccines may be used to prevent mpox disease, [JYNNEOS and ACAM2000](#).
- Clinical laboratories that perform clade-specific testing, (e.g., molecular testing or genetic sequencing) should alert and CDC ([poxvirus@cdc.gov](mailto:poxvirus@cdc.gov)) if results from such tests indicate detection of clade I MPXV.
  - Laboratories should be aware of potential genetic mutation impacts on the molecular test(s) that they are using. For instance, the subclade Ib is not detected with the previously developed "[clade I PCR test](#)". This test is now considered a clade Ia test. Visit [Lab Advisory: Recommendations for Mpox Specimen Testing](#) for additional information.
- As with all procedures, laboratories should perform a site-specific and activity-specific risk assessment to [identify and mitigate risks](#).
  - Follow U.S. CDC guidance on [infection prevention and control](#) for mpox to minimize risk when working with suspected mpox specimens.
- Contact U.S. CDC or Maine CDC for help with [specimen submission](#) to Maine's Health and Environmental Testing Laboratory.
- Specimens that cannot be accepted at U.S. CDC for clinical testing [under Clinical Laboratory Improvement Amendments \(CLIA\)](#) will be redirected for surveillance purposes and tested, providing critical data on MPXV clade(s) circulating in the United States.
- [Laboratory Response Network](#) laboratories and commercial laboratories using U.S. CDC's non-variola *orthopoxvirus* (NVO) polymerase chain reaction (PCR) test should continue submitting duplicate specimens to U.S. CDC results for routine MPXV clade-specific testing from all patients with positive NVO PCR tests.
- Be aware that specimens identified as containing a select agent, including clade unidentified or clade I MPXV, must be handled and reported in accordance with [select agent regulations](#).

## Recommendations for the General Public, Including Travelers

- Learn more about which activities may [increase your risk of exposure](#) when you travel to a country where [clade I MPXV is spreading](#). Mpox is not spread through casual contact, such as someone might have in public spaces like markets, offices, classrooms, public transit, or air travel.
- [Protect yourself and others from mpox](#), including by:
  - Avoiding close contact with people who are sick with [signs and symptoms of mpox](#), including skin or genital lesions.
  - Avoiding contact with contaminated materials, such as materials used by people who are sick (e.g., clothing, bedding, toothbrushes, or sex toys), materials used in healthcare settings, or materials that came into contact with wild animals.
  - Avoid contact with animals that can carry the virus that causes mpox or their products (e.g., bushmeat, lotions, hides) in areas where mpox is endemic, particularly in Central or West Africa.
- If you may be at risk for mpox, talk to your healthcare provider about mpox prevention, including getting vaccinated with two doses of JYNNEOS if you are [eligible to get mpox vaccine](#).
- Learn more about [Preventing Mpox While Traveling](#).
- Learn more about the current situation about [Mpox in the United States and Around the World](#).

## For More Information

### For Clinicians and Public Health Partners

- [Clade I Mpox Outbreak Originating in Central Africa | Mpox | U.S. CDC](#)
- [Ongoing Clade II Mpox Global Outbreak | Mpox | U.S. CDC](#)
- [Clinical Overview of Mpox | Mpox | U.S. CDC](#)
- [Public Health Strategies for Mpox | Mpox | U.S. CDC](#)
- [Guide to Taking a Sexual History | U.S. CDC](#)
- [Mpox Considerations for People Who Are Pregnant or Breastfeeding | Mpox | U.S. CDC](#)
- [Provider Briefing on Mpox Clade I | Mpox Briefing for Providers Who Care for Pediatric Populations | HHS](#)
- [Select Agent Regulations | Biosafety Laboratory Guidance for Handling and Processing Mpox Specimens | U.S. CDC](#)
- [Information for Clinical/Diagnostic Laboratories, Healthcare Facilities, and Other Entities Not Registered with the Federal Select Agent Program | FSAP](#)
- U.S. CDC Poxvirus and Rabies Branch: [poxvirus@cdc.gov](mailto:poxvirus@cdc.gov) or, for emergencies, U.S. CDC's 24/7 Emergency Operations Center (EOC): 770-488-7100

### For the Public

- [Mpox in the United States and Around the World: Current Situation | Mpox | U.S. CDC](#)
- [About Mpox | Mpox | U.S. CDC](#)
- [Mpox Vaccination | Mpox | U.S. CDC](#)
- [Preventing Mpox | Mpox | U.S. CDC](#)
- [Preventing Mpox While Traveling | Mpox | U.S. CDC](#)
- [Clade I Mpox in Central and Eastern Africa | September 2024 Travel Health Notice | U.S. CDC](#)
- General inquiries: U.S. CDC-INFO (1-800-232-4636)

## References

1. World Health Organization. 2022-24 Mpox (Monkeypox) Outbreak: Global Trends. [https://worldhealthorg.shinyapps.io/mpox\\_global](https://worldhealthorg.shinyapps.io/mpox_global) Accessed November 17, 2024.
2. Rao AK. "Use of JYNNEOS During Mpox Outbreaks: Clinical Guidance." Advisory Committee on Immunization Practices (ACIP) presentation. Atlanta, GA, June 23, 2023. <https://www.cdc.gov/acip/downloads/slides-2023-06-21-23/03-mpox-Rao-508.pdf>
3. Rao AK. "Evidence to Recommendations Framework: Vaccination with JYNNEOS During Mpox Outbreaks." Advisory Committee on Immunization Practices (ACIP) presentation. Atlanta, GA, February 22, 2023. <https://www.cdc.gov/acip/downloads/slides-2023-02-22-24/Mpox-07-Raohttps://www.cdc.gov/acip/downloads/slides-2023-02-22-24/Mpox-07-Rao-508.pdf508.pdf>
4. Kibungu, E. M., Vakaniaki, E. H., Kinganda-Lusamaki, E., Kalonji-Mukendi, T., Pukuta, E., Hoff, N. A., Lushima, R. S. (2024). Clade I–Associated Mpox Cases Associated with Sexual Contact, the Democratic

Republic of the Congo. *Emerging Infectious Diseases*, 30(1), 172-176.

<https://doi.org/10.3201/eid3001.231164>.

5. Yinda CK, Koukouikila-Koussounda F, Mayengue PI, et al. Genetic sequencing analysis of monkeypox virus clade I in Republic of the Congo: a cross-sectional, descriptive study. *Lancet*. 2024; 404:1815-1822. [https://www.thelancet.com/journals/lancet/article/PIIS01406736\(24\)02188-3/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS01406736(24)02188-3/fulltext)[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(24\)02188-3/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)02188-3/fulltext)