


Substance Abuse Trends in Maine

Epidemiological Profile 2013

Aroostook District



Department of Health
and Human Services

*Maine People Living
Safe, Healthy and Productive Lives*

Paul R. LePage, Governor

Mary C. Mayhew, Commissioner

Produced for Maine Department of Health and Human Services
Office of Substance Abuse and Mental Health Services
by Hornby Zeller Associates, Inc.
July 2013

Substance Abuse Trends in Maine

Epidemiological Profile 2013

Aroostook District

**THIS REPORT IS PRODUCED FOR
THE MAINE OFFICE OF SUBSTANCE ABUSE
COMMUNITY EPIDEMIOLOGY SURVEILLANCE NETWORK
WITH SUPPORT FROM THE PARTNERSHIPS FOR SUCCESS GRANT
THROUGH THE U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
SUBSTANCE ABUSE AND MENTAL HEALTH ADMINISTRATION**

**Produced by
Hornby Zeller Associates, Inc.
373 Broadway
South Portland, ME 04106
July 2013
(207) 773-9529
www.hornbyzeller.com**

Table of Contents

List of Figures	ii
Introduction	1
Overview of Aroostook Public Health District	1
Purpose of this Report	1
Consumption of Substances.....	3
Alcohol	5
Tobacco	11
Prescription Drugs.....	14
Other Illegal Drugs	16
Consequences Resulting from Substance Use and Abuse.....	19
Criminal Justice Involvement	21
Driving Under the Influence.....	30
Hospital Visits Related to Substance Use.....	34
Overdoses and Related Deaths	40
Factors Contributing to Substance Use and Abuse.....	45
Availability and Accessibility.....	47
Perceived Risk and Harm	54
Perceived Enforcement	59
Mental Health, Suicide and Co-occurring Disorders.....	61
Depression and Anxiety.....	63
Suicide and Suicidal Ideation	65
Mental Health and Substance Abuse Co-Occurrence	66
Treatment Admissions for Substance Abuse	67
Treatment Admissions.....	69
Appendix: Data Sources.....	75
Description of Data Sources	75

List of Figures

Figure 1. Percent of high school students in Aroostook who had at least one drink of alcohol during past 30 days: 2009, 2011.....	5
Figure 2. Percent of high school students in Aroostook who had at least five drinks in a row during past 30 days: 2009, 2011.....	6
Figure 3. Percent of adults by Public Health District who reported drinking during past 30 days: 2011	7
Figure 4. Percent of adults by Public Health District who reported heavy drinking during past 30 days: 2011	8
Figure 5. Percent of adults by Public Health District who reported binge drinking during past 30 days: 2011	9
Figure 6. Percent of adults by Public Health District who reported binge drinking in past 30 days by age group: 2011	10
Figure 7. Percent of high school students in Aroostook who reported smoking one or more cigarettes during past 30 days: 2009, 2011.....	11
Figure 8. Percent of high school students in Aroostook who used smokeless tobacco in the past 30 days: 2009, 2011.....	12
Figure 9. Percent of adults by Public Health District who reported smoking a cigarette in the past 30 days: 2011	13
Figure 10. Percent of high school students in Aroostook who have taken prescription drugs not prescribed to them by a doctor: 2009, 2011.....	14
Figure 11. Percent of population 12 years old or older who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2008-10	15
Figure 12. Percent of high school students in Aroostook who have used marijuana during past 30 days: 2009, 2011.....	16
Figure 13. Percent of adults who have used marijuana during the past 30 days, by Public Health District: 2011.....	17
Figure 14. Percent of high school students in Aroostook that have used cocaine in any form during their lifetime: 2009, 2011	18
Figure 15. Violent crime rate per 10,000, by Public Health District: 2006-2011.....	22
Figure 16. Alcohol-related arrest rate per 10,000, by Public Health District: 2006-2011.....	24
Figure 17. Drug-related arrest rate per 10,000, by Public Health District: 2006-2011	26

Figure 18. Drug offense arrests per 10,000 residents, by Public Health District: 2007-2012.....	28
Figure 19. Drug offense arrests per 10,000 residents in Aroostook, by drug type: 2009-12.....	29
Figure 20. Percent of high school students in Aroostook who reported drinking and driving during the past 30 days: 2009, 2011	30
Figure 21. Percent of high school students in Aroostook who rode in a vehicle driven by someone who had taken illegal drugs: 2009, 2011.....	31
Figure 22. Alcohol/Drug-related motor vehicle crash rate per 10,000, by Public Health District: 2008-2012	33
Figure 23. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District: 2010, 2011.....	35
Figure 24. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011.....	36
Figure 25. Outpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District: 2010, 2011.....	38
Figure 26. Outpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011.....	39
Figure 27. Number of overdoses per 10,000 residents, by Public Health District: 2011 and 2012	41
Figure 28. Number of overdoses per 10,000 residents, by Public Health District and primary type of substance involved: 2012.....	42
Figure 29. Drug-related death rate per 100,000, by Public Health District:	44
Figure 30. Percent of high school students in Aroostook who reported it was easy to get alcohol: 2009, 2011.....	47
Figure 31. Percent of high school students in Aroostook who reported it would be easy to get marijuana: 2009, 2011	48
Figure 32. Number of liquor licensees per 1,000 residents, by Public Health District: 2013.....	49
Figure 33. Dispensed quantity of narcotics per capita, by Public Health District: 2008-2012.....	51
Figure 34. Dispensed quantity of stimulants per capita, by Public Health District: 2008-2012.....	52

Figure 35. Dispensed quantity of tranquilizers per capita, by Public Health District: 2008-2012	53
Figure 36. Percent of high school students in Aroostook who reported a risk of harm from consuming five or more drinks once or twice per week: 2009, 2011	54
Figure 37. Percent of population age 12 or older who perceive a great risk from binge drinking, by Public Health District: 2008-10.....	55
Figure 38. Percent of high school students in Aroostook who reported a risk of harm from smoking marijuana regularly: 2009, 2011	56
Figure 39. Percent of population age 12 or older who perceive a great risk from smoking marijuana once a month by Public Health District: 2006-2008 and 2008-10	57
Figure 40. Percent of high school students who reported a risk of harm from misusing prescription drugs, by Public Health District: 2009, 2011	58
Figure 41. Perceived risk among high school students in Aroostook of being caught by parents or police for drinking alcohol: 2009, 2011	59
Figure 42. Perceived risk among high school students in Aroostook of being caught by police for smoking marijuana: 2009, 2011	60
Figure 43. Percent of adults who have ever been told they have an anxiety or depressive disorder, by Public Health District: 2011	63
Figure 44. Felt sad or hopeless almost every day for two weeks or more in a row during the past year, by Public Health District: 2011	64
Figure 45. Percent of high school students who considered suicide during the past year, by Public Health District: 2011.....	65
Figure 46. Percent of individuals in Aroostook admitted for substance abuse treatment that also had a mental health diagnosis: 2007-2012	66
Figure 47. Primary drug admissions for in Aroostook, by drug type: 2008-2012	69
Figure 48. Primary treatment admissions in Aroostook: 2012.....	70
Figure 49. Primary drug admissions per 10,000 residents, by Public Health District and drug type: 2012	71
Figure 50. Primary drug admissions per 10,000 residents, by Public Health District and drug type: 2012	72
Figure 51. Secondary treatment admissions in Aroostook: 2008-2012	73
Figure 52. Secondary treatment admissions in Aroostook, by drug type: 2012.....	74

Introduction

Overview of Aroostook Public Health District

Aroostook Public Health District has a population of 70,868 people, representing approximately 5.3 percent of Maine's total population in 2012. There are 10.8 people per square mile, making it the most rural public health district in Maine. The State of Maine is considered an "aging" state, with 16 percent of the population being 65 years old and over, a higher rate than the overall US population (13%). In Aroostook Public Health District, about 20 percent of the population was 65 years old or older in 2010. Approximately 96 percent of Aroostook's population is Caucasian, followed by African American (0.7%), American Indian and Alaska Native (1.7%), and Asian (0.5%). With a median income of \$37,138 (lowest public health district in the state), approximately 16 percent of the population lives below the poverty level. In sum, Aroostook tends to be older, poorer, and more rural compared to the rest of the state.

It is within the context of these demographic characteristics that substance abuse in Aroostook Public Health District (PHD) must be examined.

Purpose of this Report

This report takes into account the primary objectives of the Office of Substance Abuse and Mental Health Services (SAMHS): to identify substance abuse patterns in defined geographical areas, establish substance abuse trends, detect emerging substances, and provide information for policy development and program planning. It also highlights all the prevention priorities identified in the SAMHS strategic plan: underage drinking, high-risk drinking, misuse of prescription drugs, and marijuana use. Finally, the report monitors many of the factors that contribute to substance use, such as access and perceptions of harm, as well as the common negative consequences such as crime, car crashes and overdose deaths.

This report includes data available through May 2013. Older and unchanged data are included when more recent data were not available. Five major types of indicators are included: self-reported substance consumption, consequences of substance use, factors contributing to substance use, indicators about mental health and substance abuse, and treatment admissions.

The most recent data available for the Behavioral Risk Factor Surveillance System (BRFSS) are from 2011. **Due to methodological changes in weighting and sampling, 2011 BRFSS data cannot be trended with previous BRFSS years.** For this reason, we have only included snapshots of 2011 BRFSS data for this year's SEOW report.

Previous county-level reports with older trending data are available at the <http://www.maine.gov/dhhs/samhs/> website.

Consumption of Substances

Consuming harmful substances can have detrimental effects on an individual's well-being, including increased risks of morbidity, addiction and mortality, and has a harmful effect on society as a whole including increased motor vehicle accidents and crime. However, it is the manner and frequency with which people drink, smoke and use drugs that are often linked to particular substance-related consequences. To understand fully the magnitude of substance use consequences, it is important to first understand the prevalence of substance use consumption, itself. Consumption includes overall use of substances, acute or heavy consumption and consumption by high risk groups (e.g., youth, college students, pregnant women).

As demonstrated by the indicators below, alcohol remains the substance most often used by Aroostook residents across the lifespan. In particular, high-risk drinking among the youth and younger adults continues to be a concern although it appears that young people in Aroostook are somewhat less likely to engage in high-risk drinking compared to the rest of the state. Tobacco use, particularly smoking cigarettes, appears to be higher among Aroostook's population when compared to the rest of the state. Prescription drugs and marijuana are the two most commonly used drugs in Maine and Aroostook is no different, although the rates of use tend to be somewhat lower.

Alcohol


Indicator Description: ALCOHOL USE AMONG YOUTH. This measure shows the percentage of Maine high school students who reported having had one or more alcoholic drinks within 30 days prior to the survey.

Why Indicator is Important: Alcohol is the most often used substance among youth in Maine. In addition to the risks alcohol consumption carries for adults, developing adolescent brains are especially susceptible to the health risks of alcohol consumption. Adolescents who consume alcohol are more likely to have poor grades and be at risk for experiencing social problems, depression, suicidal thoughts, assault, and violence.

Data Source(s): MIYHS, 2009-2011

Summary: In Aroostook, the percentage of students who reported drinking in the past 30 days fell from 32 percent in 2009 to 27 percent in 2011, as compared to the percentage statewide, which decreased from 32 percent in 2009 to 28 percent in 2011.

Figure 1. Percent of high school students in Aroostook who had at least one drink of alcohol during past 30 days: 2009, 2011



Source: MIYHS


Indicator Description: HIGH-RISK ALCOHOL USE AMONG YOUTH. This indicator presents the percentage of Maine high school students who reported having had five or more alcoholic drinks in a row in one sitting at least once during the 30 days prior to the survey.

Why Indicator is Important: Youth are more likely than adults to engage in high-risk drinking when they consume alcohol. High-risk alcohol use contributes to violence and motor vehicle crashes and can result in negative health consequences for the consumer, including injuries and chronic liver disease. Youth who engage in high-risk drinking also are more likely to use drugs and engage in risky and antisocial behavior.

Data Source(s): MIYHS, 2009-2011

Summary: From 2009 to 2011, the percentage of high school students in Aroostook who reported having consumed five or more alcoholic beverages in one sitting during the past 30 days decreased from 19 percent to 16 percent. This is slightly less than the statewide average (17%).

Figure 2. Percent of high school students in Aroostook who had at least five drinks in a row during past 30 days: 2009, 2011



Source: MIYHS


Indicator Description: ALCOHOL USE AMONG ADULTS. This indicator portrays the percentage of adults who reported having consumed one or more alcoholic drinks on one or more days within the past 30 days.

Why Indicator is Important: Alcohol is the most often used substance in Maine adults. Excessive and high-risk alcohol use may contribute to violence and result in many negative health consequences for the consumer. Moderate drinking can also have negative health effects and lead to such consequences as alcohol-related motor vehicle crashes and increased injuries. Current alcohol use in pregnant women is also linked to low birth weight babies, sudden infant death, and other developmental delays in children.

Data Source(s): BRFSS, 2011.

Summary: In 2011, half of the adults in Aroostook reported drinking at least one alcoholic beverage within the past 30 days. This was ten percentage points lower than the statewide average and the lowest rate among public health districts.

Figure 3. Percent of adults by Public Health District who reported drinking during past 30 days: 2011



Source: BRFSS


Indicator Description: AT RISK FROM HEAVY ALCOHOL USE. This indicator examines the percentage of Maine residents who are at risk from heavy drinking in the past month. Heavy drinking is defined as two drinks per day for a man or one drink per day for a woman.

Why Indicator is Important: Heavy drinking increases the risk for many health and social related consequences. People who consume alcohol heavily are at increased risk for a variety of negative health consequences, including alcohol abuse and dependence, liver disease, certain cancers, pancreatitis, heart disease, and death. It has also been found that the more heavily a person drinks the greater the potential for problems at home, work, and with friends.¹

Data Source(s): BRFSS, 2011.

Summary: In 2011, six percent of adults in Aroostook indicated to be at risk from heavy alcohol use (1-2 drinks per day). This rate was slightly lower than the statewide average (7%) and among the lowest of the public health districts.

Figure 4. Percent of adults by Public Health District who reported heavy drinking during past 30 days: 2011



Source: BRFSS

¹ Citation from Alcoholscreening.org, a service of Join Together and the Boston University School of Public Health. Retrieved from <http://www.alcoholscreening.org/Learn-More.aspx?topicID=8&articleID=26> on 5/5/2011.


Indicator Description: HIGH-RISK ALCOHOL USE AMONG ADULTS. This indicator reflects the percentage of adults who reported engaging in high-risk “binge” drinking within the past 30 days. This is defined as five or more drinks in one sitting for a male and four or more drinks in one sitting for a female.

Why Indicator is Important: Binge drinking is considered to be a type of high-risk drinking, meaning it increases the risk for many health- and social-related consequences. It has been linked to injury (such as falls, fights, and suicides), violence, crime rates, motor vehicle crashes stroke, chronic liver disease, addiction, and some types of cancer.

Data Source(s): BRFSS, 2011.

Summary: In 2011, 14 percent of adults in Aroostook indicated they engaged in binge drinking during the past 30 days. This was two percentage points lower than the statewide average (16%) and among the lowest rates of the public health districts.


Figure 5. Percent of adults by Public Health District who reported binge drinking during past 30 days: 2011



Source: BRFSS

Summary: In 2011, the highest rate of binge drinking in Aroostook was among the 26 to 35 year old population at 31 percent; this was also the second highest rate within this age group among public health districts. The age group with the second highest binge drinking rate in Aroostook was that of the 18 to 25 year olds with 27%, followed by 36 to 49 year olds (14%), and 50 and older residents (7%).

Figure 6. Percent of adults by Public Health District who reported binge drinking in past 30 days by age group: 2011



Source: BRFSS

Tobacco


Indicator Description: SMOKING AMONG YOUTH. This indicator illustrates the percentage of Maine high school students who reported smoking a cigarette on at least one occasion within 30 days prior to the survey.

Why Indicator is Important: Use of tobacco is associated with a greater risk of negative health outcomes, including cancer, cardiovascular, and chronic respiratory diseases, as well as death.

Data Source(s): MIYHS, 2009-2011.

Summary: From 2009 to 2011 the percentage of high school students in Aroostook who reported having smoked one or more cigarettes in the past 30 days decreased from 19 percent to 17 percent. This is slightly higher than the state average (16%).

Figure 7. Percent of high school students in Aroostook who reported smoking one or more cigarettes during past 30 days: 2009, 2011



Source: MIYHS


Indicator Description: SMOKELESS TOBACCO AMONG YOUTH. This indicator illustrates the percentage of Maine high school students who reported using smokeless tobacco on at least one occasion within 30 days prior to the survey.

Why Indicator is Important: Use of tobacco is associated with a greater risk of negative health outcomes, including cancer, cardiovascular, and chronic respiratory diseases, as well as death.

Data Source(s): MIYHS, 2009-2011.

Summary: The percent of high school students in Aroostook who have used smokeless tobacco in the past 30 days remained steady at five percent from 2009 to 2011. This is lower than the statewide average (5% compared to 8%, respectively).

Figure 8. Percent of high school students in Aroostook who used smokeless tobacco in the past 30 days: 2009, 2011



Source: MIYHS


Indicator Description: SMOKING AMONG ADULTS. This indicator illustrates the percentage of Maine adults who reported using cigarettes on at least one occasion within 30 days prior to the survey.

Why Indicator is Important: Smoking is associated with a greater risk of negative health outcomes, including cancer, cardiovascular and chronic respiratory diseases, as well as death.

Data Source(s): BRFSS, 2011.

Summary: In 2011, more than one quarter (27%) of adults in Aroostook indicated they had smoked a cigarette in the past 30 days; this was one of the highest rates among public health districts and four percentage points higher than the statewide average (23%).

Figure 9. Percent of adults by Public Health District who reported smoking a cigarette in the past 30 days: 2011



Source: BRFSS

Prescription Drugs


Indicator Description: MISUSE OF PRESCRIPTION DRUGS AMONG YOUTH. This indicator presents the percentage of Maine high school students who reported using prescription drugs that were not prescribed to them by a doctor within 30 days prior to the survey.

Why Indicator is Important: Young people are increasingly using available prescription drugs, including stimulants and opiates, instead of illegal drugs to get high. Abuse of prescription drugs may lead to consequences such as unintentional poisonings or overdose, automobile crashes, addiction, and increased crime.

Data Source(s): MIYHS, 2009-2011.

Summary: In 2011, five percent of high school students in Aroostook reported having taken prescription drugs not prescribed to them by a doctor one or more times in the past 30 days, a slight decrease since 2009. This rate is lower than is found statewide (5% compared to 7%).

Figure 10. Percent of high school students in Aroostook who have taken prescription drugs not prescribed to them by a doctor: 2009, 2011



Source: MIYHS


Indicator Description: NONMEDICAL USE OF PRESCRIPTION PAIN RELIEVERS AMONG MAINERS AGE 12 AND OLDER. This measure reflects the percentage of adults who reported using prescription drugs, particularly prescription pain relievers, for reasons other than their intended purpose. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

Why Indicator is Important: Mainers are increasingly using available prescription drugs, particularly pain relievers, instead of illegal drugs to get high. Abuse of prescription drugs may lead to consequences such as unintentional poisonings, overdose, dependence and increased crime.

Data Source(s): NSDUH, 2008-10.

Summary: In 2008-10, 12 percent of 18 to 25 year olds in Aroostook reported non-medical use of prescription pain relievers during the past 30 days, higher than any other age group within the district. The second highest rate in Aroostook occurred among 12 to 17 year olds at 6% which did not vary across public health districts. Although not shown, rates of non-medical pain reliever among residents 12 and older in Aroostook changed very little from 2006-08 (4.2%) to 2008-10 (4.3%).

Figure 11. Percent of population 12 years old or older who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2008-10



Source: NSDUH

Other Illegal Drugs


Indicator Description: CURRENT MARIJUANA USE. This measure shows the percentage of Maine residents who reported using marijuana in the past 30 days. This is presented for high school students and adults in Maine.

Why Indicator is Important: Marijuana can be addictive and is associated with increased risk for respiratory illnesses and memory impairment. Even occasional use can have consequences on learning and memory, muscle coordination, and mental health symptoms.

Data Source(s): MIYHS, 2009-2011; BRFSS, 2011.

Summary: In 2011, Seventeen percent of high school students in Aroostook reported having used marijuana one or more times in the past 30 days, compared to 22 percent statewide.


Figure 12. Percent of high school students in Aroostook who have used marijuana during past 30 days: 2009, 2011



Source: MIYHS

Summary: Among Aroostook adults, four percent reported using any marijuana within the past 30 days, three percent reported using marijuana recreationally, and one percent reported using marijuana for medical reasons. With the inception of medicinal marijuana in Maine, marijuana medical use rates are expected to increase over the next few years.

Figure 13. Percent of adults who have used marijuana during the past 30 days, by Public Health District: 2011



Source: BRFSS


Indicator Description: LIFETIME COCAINE USE AMONG YOUTH. This indicator illustrates the percentage of Maine high school students who used cocaine at least once in their lifetime (i.e., ever).

Why Indicator is Important: Cocaine is highly addictive. Use of cocaine is associated with adverse health effects such as cardiac events, seizures, and stroke. It also increases the risk of cognitive impairment, injury, and crime.

Data Source(s): MIYHS, 2009-2011.

Summary: The percentage of Aroostook high school students reporting that they had used cocaine (in any form) during their lifetime has remained steady at seven percent while the statewide average decreased from 10 percent to seven percent.

Figure 14. Percent of high school students in Aroostook that have used cocaine in any form during their lifetime: 2009, 2011



Source: MIYHS

Consequences Resulting from Substance Use and Abuse

Both individuals and communities suffer the consequences of substance abuse in terms of increased health care needs and criminal justice resources. While a great deal of information regarding substance use can be obtained from the data described in the previous section, information on the effects of that use on individuals and communities can be derived from what has come to be called “consequence” data. Consequences are defined as the social, economic and health problems associated with the use of alcohol and illicit drugs. Examples are things such as illnesses related to alcohol, drug overdose deaths, property and personal crimes, as well as driving accidents, poisonings and suicides that involve alcohol or drugs.

Aroostook has consistently shown lower rates of consequences related to substance abuse, in keeping with its generally lower rates of consumption. For example, violent crime rates are lower compared to the state overall, as are car crashes related to alcohol/drugs, the rate of overdose deaths, and inpatient/outpatient admissions. However, it is important to note that drug offense rates, especially those related to methamphetamine and pharmaceutical narcotics, are relatively high compared to other public health districts and have been increasing in recent years.

Criminal Justice Involvement

Indicator Description: ANNUAL VIOLENT CRIME RATE. This indicator shows the number of violent crimes reported to the police, per 10,000 people. Violent crimes include simple and aggravated assaults, sexual assaults, and robberies. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as: $\left(\frac{\# \text{ of violent crimes}}{\text{population}}\right) \times 10,000$

Why Indicator is Important: Violence is associated with alcohol, though the causal pathway is not completely understood. Drinking on the part of the victim or a perpetrator can increase the risk of assaults and assault-related injuries. Approximately 23 percent of sexual assaults and 30 percent of physical assaults are attributable to alcohol. Reported violent crimes are an under-report of the total number of actual violent crimes.

Data Source(s): DPS, UCR, 2006-2011.

Summary: In 2011, there were 8.3 violent crimes per 10,000 people in Aroostook, compared to 12.3 per 10,000 people statewide. Although Aroostook has consistently shown lower violent crime rates compared to the state overall, the violent crime rate has been increasing in recent years.

Figure 15. Violent crime rate per 10,000, by Public Health District:
2006-2011



Source: DPS; UCR

Indicator Description: ANNUAL ALCOHOL-RELATED ARREST RATE. This indicator reflects arrests related to alcohol per 10,000 people. Alcohol-related arrests include Operating Under the Influence (OUI), liquor law violations, and drunkenness. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as: $\left(\frac{\# \text{ of alcohol arrests}}{\text{population}}\right) \times 10,000$

Why Indicator is Important: OUI and liquor law arrest rates can be an indication of the rate of criminal behavior, but it is important to note that they are also an *indication of the level of law enforcement*. Arrest rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed. The educational component of Maine's Driver Education and Evaluation Program serviced 5,500 Maine residents in 2012.

Data Source(s): DPS, UCR, 2006-2011.

Summary: In 2011, Aroostook had 60.4 alcohol-related arrests per 10,000 people, compared to the statewide rate of 73.5 per 10,000. Aroostook held the second lowest alcohol-related arrest rate among public health districts in 2011. Alcohol-related arrest rates have been steadily declining for both Aroostook and Maine since 2009.

Figure 16. Alcohol-related arrest rate per 10,000, by Public Health District:
2006-2011



Source: DPS; UCR

Indicator Description: ANNUAL DRUG-RELATED ARREST RATE. This indicator reflects the number of arrests that were related to drugs per 10,000 people. Drug-related arrests include manufacturing, sales, and possession. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time as well as make relative comparisons between small and large population areas.


Operationalized as: $\left(\frac{\# \text{ of drug arrests}}{\text{population}}\right) \times 10,000$

Why Indicator is Important: Arrest rates for drug sales, manufacturing and drug possession can be an indication of the rate of criminal behavior, but it is important to note that they are also an *indication of the level of law enforcement*. Arrests rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed.

Data Source(s): DPS, UCR, 2006-2011.

Summary: In 2011, there were approximately 46 drug-related arrests per 10,000 people in Aroostook, compared to 42.4 per 10,000 people statewide. Rates in Aroostook have been steadily increasing since 2008.

Figure 17. Drug-related arrest rate per 10,000, by Public Health District:
2006-2011



Source: DPS; UCR


Indicator Description: DRUG OFFENSE ARRESTS BY TYPE. This indicator reflects drug offense arrests made by the Maine’s Drug Enforcement Agency, overall and by drug type. The MDEA, through its eight regional multi-jurisdictional task forces, is the lead state agency in confronting drug trafficking crime. This indicator differs from the previous drug-related arrest data in that it only tracks MDEA efforts and does not encompass all activity within Maine law enforcement agencies.

Why Indicator is Important: Drug offense arrest rates can be an indication of the rate of criminal behavior, but it is important to note that they are also an indication of the level of law enforcement. Drug arrest rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed.

Data Source(s): MDEA-UCR, 2007-2012


Summary: In 2011, there were 11 drug offense arrests per 10,000 residents in Aroostook; the highest rate among public health districts in Maine. Rates have remained comparatively high since 2007. In 2012, most drug offense arrests in Aroostook were related to pharmaceutical narcotics (10 arrests per 10,000) followed by methamphetamine (9.3 arrests per 10,000).

Figure 18. Drug offense arrests per 10,000 residents, by Public Health District: 2007-2012



Source: MDEA-UCR

Figure 19. Drug offense arrests per 10,000 residents in Aroostook, by drug type: 2009-12



Source: MDEA-UCR

Driving Under the Influence


Indicator Description: DRINKING AND DRIVING AMONG YOUTH. This measure shows the proportion of high school students who reported that they drove a car after consuming alcohol at least once within 30 days prior to taking the survey.

Why Indicator is Important: Operating a vehicle after consuming alcohol increases the risk of motor vehicle crashes, injuries and death.

Data Source(s): MIYHS, 2009-2011.

Summary: In 2011, Six percent of high school students in Aroostook reported driving a vehicle at least once after drinking alcohol in the past 30 days compared to five percent statewide.

Figure 20. Percent of high school students in Aroostook who reported drinking and driving during the past 30 days: 2009, 2011



Source: MIYHS


Indicator Description: YOUTH AS PASSENGERS IN VEHICLES DRIVEN BY INDIVIDUALS USING ILLEGAL DRUGS. This measure shows the proportion of high school students who reported that within 30 days prior to taking the survey they were a passenger in a car being operated by an individual who had consumed illegal drugs.

Why Indicator is Important: Operating a vehicle while under the influence of drugs increases the risk of motor vehicle crashes, injuries and death.

Data Source(s): MIYHS, 2009-2011.

Summary: In 2011, 19 percent of high school students in Aroostook reported that, within the past 30 days, they had been passengers in a vehicle operated by someone who had taken illegal drugs, compared to 22 percent across the state.

Figure 21. Percent of high school students in Aroostook who rode in a vehicle driven by someone who had taken illegal drugs: 2009, 2011



Source: MIYHS

Indicator Description: ALCOHOL/DRUG-INVOLVED MOTOR VEHICLE CRASH RATE. This indicator shows the number of motor vehicle crashes in which alcohol or drugs were a factor per 10,000 people. Due to new data collection regulations, crash rate data is no longer separated by alcohol and drugs. Alcohol and drugs are now combined into one rate. Alcohol/drug-involved crashes means that at least one driver had consumed alcohol or drugs prior to the crash. The rate per 10,000 allows us to see the frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.

Operationalized as: $\left(\frac{\# \text{ of alcohol/drug-involved crashes}}{\text{population}}\right) \times 10,000$


Why Indicator is Important: Motor vehicle crashes are the second-leading cause of traumatic brain injury, with 27 percent of traumatic brain injuries occurring from motor vehicle crashes.² In the 2009, alcohol was attributed to 96 percent of the alcohol/drug-related crashes statewide.

Data Source(s): MDOT/MBHS, 2008-2012.

Summary: In Maine and in Aroostook, the rates of alcohol-related crashes have declined since 2008. In 2011, there were 6.6 alcohol-related crashes per 10,000 people in Aroostook; this was the lowest rate among public health districts.

² 2007 Maine Injury Report, Maine Center for Disease Control, Injury Prevention Program. Retrieved 5/17/2012 from <http://www.maine.gov/dhhs/mecdc/population-health/inj/documents/2007maineinjuryreport.pdf>

Figure 22. Alcohol/Drug-related motor vehicle crash rate per 10,000, by Public Health District: 2008-2012



Source: MDOT/MBHS

Hospital Visits Related to Substance Use

Indicator Description: INPATIENT ADMISSIONS RELATED TO SUBSTANCE USE. This indicator shows the number of inpatient hospital admissions (per 10,000 people) where alcohol, opiates, or other drugs were recorded as the primary diagnosis for which services were sought at admission. “Inpatient” refers to a patient whose treatment needs at least one night's residence in a hospital. The substance for which treatment was received was identified through hospital codes (ICD-9 codes) and includes those related to alcohol and psychoactive substances (303-305). More than one substance may be involved in a single visit. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as: $\left(\frac{\# \text{ of inpatient hospitalizations}}{\text{population}}\right) \times 10,000$

Why Indicator is Important: Hospital admissions related to substance use are an indication of injury sustained through substance use and the impact it has on the healthcare system.

Data Source(s): MHDO, 2010 and 2011.

Summary: The inpatient admissions rate due to substance use in Aroostook decreased slightly from 2010 (67.3 admissions per 10,000 residents) to 2011 (66.3 admissions per 10,000). In 2011, Aroostook had a lower inpatient substance use rate than the state (95.3 admissions per 10,000) and had the lowest rate among public health districts.

Figure 23. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District: 2010, 2011




Source: MHDO

*Visits may involve multiple substances

Summary: In 2011, most inpatient admissions due to substance use in Aroostook were related to alcohol (39.4 admissions per 10,000), followed by marijuana (14.7 admissions per 10,000), opiates (13 admissions per 10,000), and other drugs (17.4 admissions per 10,000). Among public health districts, Aroostook held the lowest inpatient rate due to alcohol as well as the lowest inpatient rate related to opiates.

Figure 24. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011



Source: MHDO

*Visits may involve multiple substances

Indicator Description: OUTPATIENT HOSPITAL VISITS RELATED TO SUBSTANCE USE. This indicator shows the number of outpatient hospital admissions (per 10,000 people) where alcohol, opiates, or other drugs was recorded as the primary diagnosis for which services were received. “Outpatient” refers to patients who receive treatment at a hospital or clinic but are not admitted overnight. The substance for which treatment was received was identified through hospital codes (ICD-9 codes) and includes those related to alcohol psychoactive substances (303-305). The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as: $\left(\frac{\# \text{ of outpatient hospitalizations}}{\text{population}}\right) \times 10,000$

Why Indicator is Important: Outpatient hospital visits related to substance use are an indication of injury sustained through substance use and the impact it has on the healthcare system.

Data Source(s): MHDO, 2010 and 2011

Summary: From 2010 to 2011, the outpatient admission rate due to substance use in Aroostook decreased slightly from 340.3 admissions per 10,000 residents to 284.6 admissions per 10,000 residents. In 2011, Aroostook had a much lower rate than the state (708.7 admissions per 10,000) and held the lowest rate among public health districts.

Figure 25. Outpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District: 2010, 2011




Source: MHDO

*Visits may involve multiple substances

Summary: In 2011, Aroostook had the second lowest alcohol-related outpatient admission rate among public health districts (559.9 admissions per 10,000); this rate was lower than that of the state (291.1 admissions per 10,000). Aroostook also held the lowest opiate-related outpatient rate at 48 admissions per 10,000 residents; this was much lower than the statewide rate (126.4 admissions per 10,000).

Figure 26. Outpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011



Source: MHDO

*Visits may involve multiple substances

**Includes prescription narcotics, methadone, and heroin.

Overdoses and Related Deaths

Indicator Description: OVERDOSES. This indicator shows the number of persons receiving help from Emergency Medical Services (EMS) related to an overdose in 2011 and 2012.


Why Indicator is Important: Overdosing on a substance can cause serious physical harm resulting in hospitalization and even death. Responding to overdoses also uses valuable EMS resources. The rate per 10,000 allows us to see the frequency with which an occurrence happens within a population over time, as well as make relative comparisons between small and large population areas. In this case, the base of 10,000 people was used due to small numbers.

Operationalized as: $\left(\frac{\# \text{ of overdose deaths}}{\text{population}}\right) \times 10,000$

Data Source(s): Emergency Medical Services, 2011-2012


Summary: From 2011 to 2012, the Aroostook EMS overdose rate increased from 23.8 overdoses per 10,000 residents to 27.4 overdoses per 10,000 residents. In 2012, Aroostook had the second lowest EMS overdose rate among public health districts. When examined by primary type of substance involved, alcohol had the highest rate with 19.2 overdoses per 10,000 residents followed by medication with 17.9 overdoses per 10,000 residents.

Figure 27. Number of overdoses per 10,000 residents, by Public Health District: 2011 and 2012



Source: Emergency Medical Services

Figure 28. Number of overdoses per 10,000 residents, by Public Health District and primary type of substance involved: 2012



Source: Emergency Medical Services, 2012

Indicator Description: DEATHS DUE TO OVERDOSE. This measure reflects the number of deaths where the cause of death was directly related to the consumption of one or more substances. The measure excludes deaths where a substance may have been ingested prior to engaging in a behavior that resulted in death (e.g., drunk driving) or where lifetime substance use and abuse may have impacted health (e.g., cirrhosis). In order to preserve anonymity and strengthen validity, rates were calculated based on the sum of deaths per three year intervals. The rate per 100,000 allows us to see the frequency with which an occurrence happens within a population over time, as well as make relative comparisons between small and large population areas. In this case, the base of 100,000 people was used due to small numbers.


Operationalized as: $\left(\frac{\# \text{ of overdose deaths}}{\text{population}}\right) \times 100,000$

Why Indicator is Important: One of the most extreme consequences of alcohol and drug abuse is overdose death; that is, the substance(s) consumed played a direct role in an individual's death. These are seen as potentially preventable deaths.

Data Source(s): Office of Chief Medical Examiner, 2000-02 to 2009-11.

Summary: Between years 2009 and 2011, the drug overdose death rate in Aroostook was 28 per 100,000; this rate was the lowest among public health districts but has been increasing since the period of 2000-02 (8.2 per 10,000).

Figure 29. Drug-related death rate per 100,000, by Public Health District:
2000-02 to 2009-11



Source: Office of the Chief Medical Examiner.

Factors Contributing to Substance Use and Abuse

A body of substance abuse prevention research has identified certain groups of factors that “cause” or have an impact on substance use and the consequences related to use. That is, they appear to influence the occurrence and magnitude of substance use and its related consequences. Generically, these causal factors (also known as contributing factors) are categorized into groups which include:

- Social Access (e.g., getting drugs and alcohol from friends or family)
- Retail Availability (e.g., retailer not carding properly)
- Pricing & Promotion (e.g., two-for-one specials, industry sponsorships or signage)
- Social/Community Norms (e.g., parental/community attitudes and beliefs)
- Enforcement (e.g., lack of compliance checks)
- Perceptions of Harm (e.g., individuals’ belief that using a substance is harmful)³
- Perceived Risk of Being Caught (e.g., individuals’ belief that s/he will be caught by parents or police)⁴

Substance abuse prevention in Maine is undertaken with the assumption that making changes to these factors at the community level will result in changing behaviors around substance use and related problems. It is through positively impacting these factors that Maine can achieve population-level changes in substance consumption and consequences.

Although most high school students in Aroostook seem to perceive that regular use of substances poses a risk of harm, less than half think they will be caught by their parents and even fewer think they will be caught by the police if they use alcohol or marijuana. In fact, most students in Aroostook think it is easy to obtain alcohol and marijuana.

³ Bonnie, Richard J., and Mary Ellen O’Connell, Eds. (2004). *Reducing Underage Drinking: A Collective Responsibility*. The National Academies Press: Washington, DC.

⁴ "A General Causal Model to Guide Alcohol, Tobacco and Illicit Drug Prevention: Assessing the Research Evidence." Multi-State Technical Assistance Workshop. Washington, DC. March 16, 2006.

Availability and Accessibility


Indicator Description: PERCEIVED EASE OF OBTAINING ALCOHOL BY UNDERAGE YOUTH. This indicator reflects the percentage of high school students (grades 9 to 12) who reported that it would be easy or very easy for them to get alcohol if they wanted some.

Why Indicator is Important: According to the 2011 statewide MIYHS, students who reported that they thought alcohol was easy to obtain were three times as likely to report consuming alcohol within the past 30 days compared to students who did not think it was easy obtain.

Data Source(s): MIYHS, 2009-2011.

Summary: From 2009 to 2011, the percentage of high school students in Aroostook who indicated that it was easy to get alcohol decreased from 70 percent to 66 percent. This was slightly lower than the statewide rate of 67 percent.

Figure 30. Percent of high school students in Aroostook who reported it was easy to get alcohol: 2009, 2011



Source: MIYHS


Indicator Description: PERCEIVED EASE OF OBTAINING MARIJUANA BY YOUTH. This indicator illustrates the percentage of high school students reporting it would be easy or very easy to obtain marijuana if they wanted it.

Why Indicator is Important: According to the 2011 statewide MIYHS, students who reported that they thought marijuana was easy to obtain were seven times as likely to use marijuana in the past 30 days compared to their peers who thought it was difficult to obtain.

Data Source(s): MIYHS, 2009-2011.

Summary: In 2011, Fifty percent of high school students in Aroostook indicated that it would be easy to get marijuana; the state average was notably higher (58%).

Figure 31. Percent of high school students in Aroostook who reported it would be easy to get marijuana: 2009, 2011



Source: MIYHS


Indicator Description: NUMBER OF ALCOHOL OUTLETS PER CAPITA. This indicator reflects the number of active (as of May 2013) retail establishments selling alcohol per person. This includes both on-premise (e.g., bars, restaurants) and off-premise (e.g., convenience stores) establishments. It is calculated by dividing the number of retail establishments by the number of residents in the county (based on 2010 U.S. Census figures).

Why Indicator is Important: National research shows that there is a correlation between the number of places that sell alcohol in an area (retail density) and the rate of alcohol-related crime.⁵

Data Source(s): DPS, Liquor Licensing and Compliance, 2013; U.S. Census, 2010.

Summary: In 2013, the number of active liquor licenses in Aroostook per 1,000 residents (3.8) was higher than the statewide average (3.1) and second highest among public health districts.

Figure 32. Number of liquor licenses per 1,000 residents, by Public Health District: 2013



Source: DPS/U.S. Census

⁵ Grube, J. W., Gruenewald, P. J. & Chen, M. J. (2010). Community alcohol outlet density and underage drinking. *Addiction*, 105, 270-278.


Indicator Description: DISPENSED QUANTITY OF SCHEDULE II DRUGS PER CAPITA. These indicators reflect the dispensed quantity of narcotics, tranquilizers, and stimulants through prescriptions in Maine. This includes only prescription drugs that are classified “Schedule II” drugs, meaning those with a high potential for abuse. It is important to note that the dispensed quantity does not indicate the size or dosage of the pills associated with the prescription. All pharmacies in Maine report to the Prescription Monitoring Program.

Why Indicator is Important: The dispensed quantity per capita indicates the volume of prescription drugs potentially available in the community for diversion (e.g., gift, sale, or theft). A higher level of availability contributes to misuse by individuals without a prescription.

Data Source(s): PMP, 2008-2012

Summary: In 2012, dispensed quantity of narcotics per capita in Aroostook was the second lowest among public health districts and has been decreasing for the past five years shown (from 69 per 10,000 in 2008 to 52.8 per 10,000 in 2012).


Figure 33. Dispensed quantity of narcotics per capita, by Public Health District: 2008-2012



Source: PMP, 2006-2012

Summary: The dispensed quantity of stimulants per capita in Aroostook has steadily increased from 8.5 per person in 2008 to 9.8 per person in 2012. Aroostook has consistently held some of the lowest rates among public health districts for past several years shown.


Figure 34. Dispensed quantity of stimulants per capita, by Public Health District: 2008-2012



Source: PMP, 2008-2012

Summary: The dispensed quantity of tranquilizers per capita in Aroostook decreased slightly from 29.3 in 2008 to 28.3 in 2012. Most public health districts' rates, including Aroostook, remained relatively stable and have remained on par with the statewide average over the past five years shown.

Figure 35. Dispensed quantity of tranquilizers per capita, by Public Health District: 2008-2012



Source: PMP, 2008-2012

Perceived Risk and Harm


Indicator Description: PERCEIVED RISK FROM BINGE DRINKING AMONG YOUTH. This indicator reflects the percentage of individuals who perceive that there is moderate-to-great risk from drinking five or more drinks once or twice per week.

Why Indicator is Important: According to the 2011 statewide MIYHS, high school students who perceive binge drinking as a moderate-to-great risk of harm are one-third as likely to binge drink in the past 30 days than students who did not perceive harm. Adults are also less likely to binge drink if they perceive it to be risky.

Data Source(s): MIYHS, 2009-2011.

Summary: From 2009 to 2011, the percentage of high school students in Aroostook who indicated that there is a moderate-to-great risk of people harming themselves if they consume five or more drinks regularly increased from 75 percent to 80 percent. This is higher than the state average (78%).

Figure 36. Percent of high school students in Aroostook who reported a risk of harm from consuming five or more drinks once or twice per week: 2009, 2011



Source: MIYHS


Indicator Description: PERCEIVED RISK FROM BINGE DRINKING AMONG MAINERS. This indicator reflects the percentage of Mainers age 18 and older who perceive that there is risk from consuming five or more drinks once or twice per week. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

Why Indicator is Important: The perception that consuming a lot of alcohol is risky indicates an individual is knowledgeable about health risks and other negative consequences. Adults are less likely to binge drink if they perceive it to be risky.

Data Source(s): NSDUH, 2008-10.

Summary: During the period 2008-10, Aroostook/Downeast residents ages 18-25 indicated the lowest perception of risk from binge drinking among age groups within the district at 29 percent, while those 26 and older had the highest proportion (40%). Aroostook/Downeast’s rates of perception are very similar to those of the state. Although not shown, rates of perception of risk from binge drinking among Aroostook/Downeast residents 12 and older remained unchanged since 2006-08 (39%).

Figure 37. Percent of population age 12 or older who perceive a great risk from binge drinking, by Public Health District: 2008-10



Source: NSDUH


Indicator Description: PERCEIVED RISK OF REGULAR MARIJUANA USE AMONG YOUTH. This measure demonstrates the percentage of individuals who perceive a moderate-to-great risk of harm from smoking marijuana regularly.

Why Indicator is Important: According to the 2011 statewide MIYHS, high school students who do not believe there is moderate-to-great risk in smoking marijuana regularly are 6.5 times as likely to smoke marijuana as their peers who do perceive risk of harm.

Data Source(s): MIYHS, 2009-2011.

Summary: From 2009 to 2011, the percentage of high school students in Aroostook who indicated that there is a moderate-to-great risk of people harming themselves if they smoke marijuana regularly decreased from 70 percent to 64 percent. The percentage of Aroostook high school students holding this view was substantially greater than the state average (56%).

Figure 38. Percent of high school students in Aroostook who reported a risk of harm from smoking marijuana regularly: 2009, 2011



Source: MIYHS


Indicator Description: PERCEIVED RISK OF MARIJUANA USE AMONG MAINERS. This measure demonstrates the percentage of Mainers over the age of 12 who perceive a risk of harm from smoking marijuana once a month. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

Why Indicator is Important: The perception that using a substance is risky indicates an individual is knowledgeable about health risks and other negative consequences associated with that substance. Perceptions of risk reduce the likelihood that an individual will engage in the behavior.

Data Source(s): NSDUH, 2006-08 and 2008-10.

Summary: From 2006-08 to 2008-10, the percentage of Aroostook/Downeast residents 12 and older who perceived a great risk from smoking marijuana once a month decreased from 36 percent to 33 percent. Aroostook/Downeast had the highest perception of risk among public health districts in 2008-10.

Figure 39. Percent of population age 12 or older who perceive a great risk from smoking marijuana once a month by Public Health District: 2006-2008 and 2008-10



Source:

NSDUH


Indicator Description: PERCEIVED RISK OF PRESCRIPTION DRUG MISUSE AMONG YOUTH. This measure demonstrates the percentage of individuals who perceive a moderate-to-great risk of harm from taking a prescription drug that was not prescribed to them.

Why Indicator is Important: According to the 2011 statewide MIYHS, high school students who do not believe there is moderate-to-great risk misusing prescription drugs are 4.6 times as likely to smoke marijuana as their peers who do perceive risk of harm.

Data Source(s): MIYHS, 2009-2011.

Summary: Perception of risk from misusing prescription drugs among high school students in Aroostook remained the same from 2009 to 2011 at 87 percent; this was slightly higher than the statewide average (85%). That means 13 percent of students in Aroostook did not think misusing prescription drugs was risky.

Figure 40. Percent of high school students who reported a risk of harm from misusing prescription drugs, by Public Health District: 2009, 2011



Source: NSDUH

Perceived Enforcement


Indicator Description: PERCIEVED RISK OF BEING CAUGHT FOR DRINKING ALCOHOL AMONG YOUTH. This indicator reflects the percentage of high school students who reported that they would be caught by their parents or by police if they drank alcohol.

Why Indicator is important: According to the 2011 statewide MIYHS, high school students who believe they will be caught by their parents are one-fifth as likely to drink in the past month as compared to students who do not think they will be caught. Students who believe that they would be caught by the police are half as likely to drink alcohol in the past month as those who do not think they would be caught.

Data Source(s): MIYHS, 2009-2011.

Summary: At 47 percent, the perceived risk among high school students of being caught by their parents for drinking alcohol in Aroostook is greater than the state average (43%). Only 14 percent of high school students in Aroostook indicated that they thought they would be caught by the police for drinking alcohol (compared to 15% statewide). That means high school students in Aroostook are more than three times as likely to perceive a risk of being caught by their parents, rather than the police, for drinking alcohol.

Figure 41. Perceived risk among high school students in Aroostook of being caught by parents or police for drinking alcohol: 2009, 2011



Source: MIYHS


Indicator Description: PERCEIVED RISK OF BEING CAUGHT FOR SMOKING MARIJUANA AMONG YOUTH. This measure shows the percentage of high school students who reported that they thought they would be caught by police if they smoked marijuana.

Why Indicator is Important: According to the statewide 2011 MIYHS, high school students who believe they would be caught by the police are approximately half as likely to smoke marijuana as their peers.

Data Source(s): MIYHS, 2009-2011.

Summary: In 2011, 29 percent of high school students in Aroostook indicated that they thought they would be caught by the police if they smoked marijuana, compared to 25 percent statewide.

Figure 42. Perceived risk among high school students in Aroostook of being caught by police for smoking marijuana: 2009, 2011



Source: MIYHS

Mental Health, Suicide and Co-occurring Disorders

The relationship between substance use and mental health has been well documented. There are great efforts underway at the Substance Abuse Mental Health Services Administration (SAMHSA) and throughout Maine to better integrate mental health promotion and substance abuse prevention. At the individual level, it is important to know if one exists because the symptoms of each can affect the other; that is, a person who is depressed may abuse alcohol or drugs in an effort to feel better. At the community level, it is important to understand how the prevalence of one interacts with the other so that prevention and intervention efforts can better address the needs of both. The data indicators included below represent the first attempt to collect multiple mental health indicators that can be routinely monitored in relation to substance abuse in hopes that this will lead to better prevention and intervention.

About one-fifth of adults in Aroostook report having ever been diagnosed with depression or anxiety and about one quarter of high school students felt sad or hopeless every day for two weeks in 2011; one in ten high school students considered suicide. The proportion of individuals from Aroostook admitted for substance abuse treatment who also have a mental health diagnosis has been increasing since 2007.

Depression and Anxiety


Indicator Description: DIAGNOSIS OF ANXIETY AND DEPRESSION AMONG ADULTS. This indicator examines the percentage of Maine residents age 18 and older who have ever been told by a doctor that they have a depressive or anxiety disorder.

Why Indicator is Important: The link between mental health and substance abuse is well documented. Experiencing anxiety or depression is associated with higher rates of substance abuse.

Data Source(s): BRFSS, 2011.

Summary: In 2011, approximately 18 percent of adults in Aroostook had been told they have an anxiety disorder, and 24 percent had been told they have a depressive disorder. The rates for diagnosis of anxiety and depressive disorders in Aroostook are relatively on par with statewide rates.

Figure 43. Percent of adults who have ever been told they have an anxiety or depressive disorder, by Public Health District: 2011



Source: BRFSS


Indicator Description: DEPRESSION AMONG YOUTH. This indicator measures the percentage of high school students reporting they felt sad or hopeless almost every day for two weeks in a row during the past year.

Why Indicator is Important: Experiencing depression in the past year is associated with higher rates of substance abuse. Among youth, depression is also associated with problems with relationships and academic achievement.

Data Source(s): MIYHS, 2011.

Summary: In 2011, 23 percent of high school students in Aroostook indicated that they felt sad or hopeless every day for two weeks or more in a row during the past year. This rate was the same found amongst Maine high school students overall.

Figure 44. Felt sad or hopeless almost every day for two weeks or more in a row during the past year, by Public Health District: 2011



Source: MIYHS

Suicide and Suicidal Ideation


Indicator Description: SUICIDAL IDEATION AMONG YOUTH. This measure examines the percentage of high school students who reported that they seriously considered attempting suicide during the past year.

Why Indicator is Important: Suicide is the most tragic consequence of major depressive disorders. Abuse of alcohol or other drugs may increase emotional problems leading to suicidal ideation and suicidal behavior.

Data Source(s): MIYHS, 2011.

Summary: In 2011 the percentage of high school students who considered suicide during the past year was slightly lower in Aroostook (12%) than statewide (13%).

Figure 45. Percent of high school students who considered suicide during the past year, by Public Health District: 2011



Source: MIYHS

Mental Health and Substance Abuse Co-Occurrence


Indicator Description: CO-OCCURRING MENTAL HEALTH AND SUBSTANCE ABUSE TREATMENT. This indicator reflects the proportion of treatment admissions for substance abuse where the individual also has a mental health diagnosis.

Why Indicator is Important: The link between mental health and substance abuse is well documented. In terms of treatment, it is important to know if one exists because the symptoms of each can affect the other.

Data Source(s): TDS, 2007-2012.

Summary: From 2007 through 2012, Aroostook has consistently reported significantly lower percentages of individuals admitted for substance abuse treatment who also have a mental health diagnosis than is found in the state on average. However, Aroostook has seen an increase of eight percentage points from 33% in 2007 to 41 percent in 2012.

Figure 46. Percent of individuals in Aroostook admitted for substance abuse treatment that also had a mental health diagnosis: 2007-2012



Source: TDS

Treatment Admissions for Substance Abuse

Substance abuse treatment admissions are an indicator of how many people *receive treatment* for a substance abuse problem. These admissions can be voluntary, but they can also be court-ordered. Treatment admission data should not be used as an indicator of the magnitude of the problems related to substance abuse. Rather, treatment should be seen as a major consequence stemming from substance use and one that requires many resources.

The overall number of clients admitted to treatment has been declining since 2007, from 14,843 to 11,688 in 2012. Mainers continued to seek out treatment for abuse involving a wide array of substances besides alcohol; in 2012 there were 4,135 admissions for alcohol as the primary substance. This was followed by synthetic opioids (3,838) and marijuana (1,024).

In Aroostook, two-fifths of primary treatment admissions were for alcohol and one-third were for synthetic opioids; these rates appear to have remained relatively stable over the past several years. In 2012, among public health districts, Aroostook had the highest rate of alcohol-related primary admissions per 10,000 residents, the highest rate of marijuana-related primary admissions per 10,000 residents, and the second highest rate of methadone-related primary admissions per 10,000 residents.

Treatment Admissions


Indicator Description: PRIMARY TREATMENT ADMISSIONS. This measure reflects substance abuse treatment admissions. A “primary” substance is identified during the treatment admissions process based on use patterns (e.g., frequency, duration, quantity) and the risk(s) posed to the individual. The analysis excludes admissions for shelter/detoxification services.

Why Indicator is Important: The number of substance abuse treatment admissions is bound by both the need and the capacity for treatment. Treatment admissions data are not a good indicator of substance use, abuse or dependence, but provide an indication of service usage and the impact of substance use on the behavioral healthcare system.

Data Source(s): TDS, 2009-2012.

Summary: In 2011, 42 percent of all primary treatment admissions in Aroostook were related to alcohol, followed by synthetic opioids⁶ (29%) and marijuana (12%). This trend has remained relatively stable since 2007 although primary treatment admissions related to alcohol appear to be gradually declining as a proportion of all admissions.

Figure 47. Primary drug admissions for in Aroostook, by drug type: 2008-2012




Source: TDS

⁶ “Synthetic opioids” excludes methadone and buprenorphine.

Summary: In 2012, the proportion of primary treatment admissions for alcohol in Aroostook was six percentage points higher than the statewide average (42% compared to 36%), while the rate for synthetic opioids was four percentage points lower than the state (29% compared to 33%). At 12 percent, Aroostook had a higher percentage of primary admissions related to marijuana compared to the primary treatment admissions statewide (9%).


Figure 48. Primary treatment admissions in Aroostook: 2012



Source: TDS

Summary: In 2012, Aroostook had the highest rate among public health districts of primary treatment admissions due to alcohol (47 admissions per 10,000 residents) and the fourth highest rate of primary admissions due to synthetic opioids (33 admissions per 10,000 residents).


Figure 49. Primary drug admissions per 10,000 residents, by Public Health District and drug type: 2012



Source: TDS

Summary: In 2012, Aroostook had the highest rate among public health districts of primary admissions due to marijuana (13.5 admissions per 10,000), the second highest rate of primary admissions due to methadone (6.2 admissions per 10,000 residents), the lowest rate of primary admissions due to heroin/morphine (2.4 admissions per 10,000 residents), and the second lowest rate of primary admissions due to cocaine/crack (1.3 admissions per 10,000 residents).

Figure 50. Primary drug admissions per 10,000 residents, by Public Health District and drug type: 2012



Source: TDS


Indicator Description: SECONDARY TREATMENT ADMISSIONS. This measure reflects substance abuse treatment admissions. A “secondary” substance is identified during the admissions process as one used by the individual and for which treatment may be received, but it is not the primary substance for which treatment was sought. The analysis excludes admissions for shelter/detoxification services.

Why Indicator is Important: The number of substance abuse treatment admissions is bound by both the need and the capacity for treatment. Treatment admission data are not a good indicator of substance use, abuse or dependence, but provide an indication of service usage and the impact of substance use on the behavioral healthcare system.

Data Source(s): TDS, 2008-2012.

Summary: In 2012, 34 percent of secondary treatment admissions in Aroostook were for marijuana, followed by synthetic opioids (22%) and then alcohol (16%). These trends have remained relatively stable since 2008 although it appears that synthetic opioids as a secondary reason for treatment is decreasing.


Figure 51. Secondary treatment admissions in Aroostook: 2008-2012



Source: TDS

Summary: In 2012, Aroostook had a higher percent of secondary admissions related to marijuana (34%), alcohol (16%), and other drugs (13%) compared to secondary treatment admissions statewide.

Figure 52. Secondary treatment admissions in Aroostook, by drug type: 2012



Source: TDS

Appendix: Data Sources

This report includes data that was gathered from a number of data sources. A detailed description of each source is provided below, consisting of information about the data included in each source, and retrieval or contact information. The report includes data that were available through May 2013.

There are multiple purposes for this report. One is to provide a snapshot of the most recent data regarding substance abuse, while another is to examine trends over time. Therefore, each indicator may have multiple sources of data that are included. While each indicator provides a unique and important perspective on drug use in Maine, none should individually be interpreted as providing a full picture of drug trends in Maine. In particular, the percentages and figures from one data source do not always align with the data and percentages from a similar source. Older data are often included in order to examine an indicator among a specific population or to find trends over time. When discussing rates of prevalence, however, the user should rely upon the most recent data source available.

Description of Data Sources

Behavioral Risk Factor Surveillance System (BRFSS). The BRFSS is a national survey administered on an ongoing basis by the National Centers for Disease Control and Prevention (CDC) to adults in all 50 states and several districts and territories. The instrument collects data on adult risk behaviors, including alcohol abuse. BRFSS defines heavy drinking as adult men having more than two drinks per day and adult women having more than one drink per day, and binge drinking as males having five or more drinks on one occasion and females having four or more drinks on one occasion. The most recent data available are from 2011. **Due to methodological changes in weighting and sampling, 2011 BRFSS data cannot be trended with previous BRFSS years.** For this reason, we have only included snapshots of 2011 BRFSS data for this year's SEOW report. Both state and national data are available. Contact: Kim E. Haggan, Maine BRFSS Acting Coordinator; kim.e.haggan@maine.gov; (207) 287-5459.

Maine Department of Public Safety (DPS), Uniform Crime Reports (UCR). UCR data include drug and alcohol arrests. Drug arrests include sale and manufacturing as well as possession of illegal substances. Liquor arrests include all liquor law violations. OUI arrests are arrests for operating a motor vehicle under the influence of a controlled substance. DPS data are now available from 2011. Arrest data may reflect differences in resources or focus of law enforcement efforts, so may not be directly comparable from year to year.

Retrieval: http://www.maine.gov/dps/cim/crime_in_maine/cim.htm

Maine Department of Public Safety (DPS), Liquor Licensing and Compliance. DPS issues and renews licenses for the manufacture, importation, storage, transportation and sale of all liquor and administers those laws relating to licensing and the collection of taxes on malt liquor and wine. DPS maintains a list of all active licenses that can be accessed online.

Retrieval: http://www.maine.gov/dps/liqr/active_licenses.htm

Maine Department of Public Safety (DPS), Bureau of Highway Safety (BHS), Maine Department of Transportation (MDOT). The Bureau of Highway Safety is responsible for tracking all fatalities that occur on Maine's highways and reporting this information through the Fatal Analysis Reporting System (FARS). The data represented provides information on highway crashes and fatalities. Much of this information is gathered from our FARS system, which records data on fatal crashes in Maine for input into a larger national record-keeping system of statistical data. FARS data is also used by BHS and the Maine State Police to analyze enforcement priorities and schedules. Impaired driving is one of the most serious traffic risks facing the nation, killing thousands every year. Contact: Duane Brunell, Safety Performance Analysis Manager; duane.brunell@maine.gov; (207) 624-3278.

Maine Drug Enforcement Agency (MDEA). The MDEA, through its eight regional multi-jurisdictional task forces, is the lead state agency in confronting drug trafficking crime. This indicator differs from the Uniform Crime Report drug-related arrest data in that it only tracks MDEA efforts and does not encompass all activity within Maine law enforcement agencies. The data included in this report represents those arrested for a drug offense but does not indicate what other drug(s) may have been seized. For example, a person may be arrested for the sale of cocaine but also be in possession of oxycodone and marijuana. It is important to note that arrests and multi-jurisdictional drug enforcement are resource-dependent; such funds fluctuate from year to year, and must be reallocated to combat highest priority threats. Contact: Roy E. McKinney, Director; roy.e.mckinney@maine.gov; (207) 626-3852.

Maine Emergency Medical Services (EMS). Maine EMS is a bureau within the Maine Department of Public Safety (DPS) and is responsible for the coordination and integration of all state activities concerning Emergency Medical Services and the overall planning, evaluation, coordination, facilitation and regulation of EMS systems. EMS collects data statewide from the 272 licensed ambulance and non-transporting services. It is mandated that services submit an electronic patient care report to Maine EMS within one business day of patient contact. Data are compiled upon request. Contact: Jon Powers, Maine Emergency Medical Services; jon.powers@maine.gov; (207) 626-3860.

Maine Integrated Youth Health Survey (MIYHS). The MIYHS is a statewide survey administered biennially through a collaborative partnership by the Maine Office of Substance Abuse (OSA) the Maine Center for Disease Control and Prevention and the Maine department of Education to students in grades 5 through 12. The survey collects information on student substance use, risk factors related to substance use, as well as consequences, perceptions and social risk factors related to substances, and collects information on many other health factors. As of the date of this report, the most recent data available are from 2011. Due to changes in the survey administration and structure, the new survey data cannot be trended with the Maine Youth Drug and Alcohol Survey (MYDAUS). Contact: Stephen Corral, Substance Abuse Program Specialist, Office of Substance Abuse, stephen.corral@maine.gov; (207) 287-2964.

Maine Health Data Organization (MHDO). MHDO data includes all inpatient admissions to all hospitals in Maine for calendar years 2010 and 2011. Data categories created by the authors include alcohol, opioids, illegal drugs, and pharmaceuticals. All drug categories include intoxication, abuse, dependence, and poisoning cases related to the drug. The opiate category includes methadone, heroin, and opiates. The illegal drug category includes crack/cocaine, cannabis, and hallucinogens. The pharmaceuticals category includes all other non-opioid medications (including stimulants and depressants). Contact: Maine Health Data Organization (MHDO), lisa.parker@maine.gov; (207) 287-3225.

Maine Office of the Chief Medical Examiner. The Maine Office of the Chief Medical Examiner maintains records of all deaths associated with drug overdose. Drug categories include methadone, cocaine, benzodiazepines, oxycodone and heroin/morphine. The death data are compiled on an annual basis and must be finalized prior to release and so are not available to track changes that may occur over shorter time frames. Contact: Dr. Marcella Sorg, Director, Rural Drug & Alcohol Research Program, Margaret Chase Smith Policy Center, University of Maine; marcella_sorg@umit.maine.edu; (207) 581-2596.

National Survey on Substance Use and Health (NSDUH). The NSDUH is a national survey administered annually by the Substance Abuse and Mental Health Services Administration (SAMHSA) to youth grades 6 through 12 and adults ages 18 and up. The instrument collects information on substance use and health at the national, regional and state levels. The advantage of NSDUH is that it allows comparisons to be made across the lifespan (that is, ages 12 and up). However, NSDUH is not as current as other data sources; as of this report, data at the state level are available from 2009-2010. Older data are included for trending and comparative purposes. NSDUH defines Illicit Drugs as marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used non-medically; Binge Alcohol Use as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least one day in the past 30 days; Dependence or abuse based on definitions found in the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV); and Serious Mental Illness (SMI) as a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. Retrieval: <https://nsduhweb.rti.org/>

Northern New England Poison Center (NNEPC). The Northern New England Poison Center provides services to Maine, New Hampshire, and Vermont. A poisoning case represents a single individual's contact with a potentially toxic substance. Intentional poisoning includes those related to substance abuse, suicide and misuse. Data include the number of confirmed cases where exposures are judged to be substance abuse-related (i.e., an individual's attempt to get high). NNEPC collects detailed data on specific substances involved in poisonings, including the categories of stimulants/street drugs, alcohol, opioids, asthma/cold and cough, benzodiazepines, antidepressants, and pharmaceuticals, as well as other substances. The category of stimulants/street drugs includes marijuana and other cannabis, amphetamine and

amphetamine-like substances, cocaine (salt and crack), amphetamine/dextroamphetamine, caffeine tablets/capsules, ecstasy, methamphetamine, GHB, and other/unknown stimulants/street drugs. The category alcohol includes alcohol-containing products such as mouthwash. The opioid category includes Oxycodone, Hydrocodone, buprenorphine, methadone, tramadol, morphine, propoxyphene, codeine, hydromorphone, stomach opioids, Meperidine (Demerol), heroin, Fentanyl, and other/unknown opioids. The asthma/cold and cough category includes eye, ear, nose, and throat medications. Data available from the poison center are reported on a continual daily basis and are included through December 2012. These data are only reflective of cases in which the Poison Center was contacted. Contact: Karen Simone, Director, Northern New England Poison Center; simonk@mmc.org; (207) 662-7221.

Prescription Monitoring Program (PMP). PMP maintains a database of all transactions for class C-II through C-IV drugs dispensed in the state of Maine. Drug categories used in this report include narcotics, tranquilizers, stimulants, and other prescriptions. Other prescriptions include those drugs that are not classified as narcotics, tranquilizers or stimulants, including products such as endocrine and metabolic drugs, analgesics and anesthetics, gastrointestinal agents, and nutritional products. The counts included in this report represent the quantity dispensed through prescriptions filled between 2006 and 2012. Contact: John Lipovsky, PMP Coordinator, Substance Abuse and Mental Health Services; john.lipovsky@maine.gov; (207) 287-3363.

Treatment Data System (TDS). TDS is a statewide database that includes information about clients admitted to treatment in OSA-funded facilities through December 2012. Analyses in this report are based on clients' reported primary, secondary and tertiary drug(s) of choice as well as other demographic and background information that is collected at intake. Drug categories included in this report are alcohol, marijuana, cocaine, heroin, synthetic opiates, methadone/buprenorphine and benzodiazepines. Contact: Stacey Chandler, Substance Abuse and Mental Health Services; stacey.chandler@maine.gov; (207) 287-6337.

U.S. Census Bureau. The U.S. Census provides summary profiles showing frequently requested data items from various Census Bureau programs. Profiles are available for all states and counties, and for cities and towns with more than 25,000 people. Data are updated no less than annually. Retrieval for Maine census data: <http://quickfacts.census.gov/qfd/states/23000.html>